



“Litterfall and Leaf Area Index Measurements Before and After Selective Logging in Tapajós National Forest, Santarém - Pará - Brasil”

¹Figueira, A.M.S.; ¹Sousa, C.A.; ¹Maia, A.; ²Rocha, H.; ²Juarez, R.; ²Freitas, H.; ³Goulden, M.; ³Menton, M.; ³Miller, S.D.

¹Universidade Federal do Pará – Campus de Santarém-PA, ²Universidade de São Paulo, ³University of California, Irvine

Introduction

Biophysical processes and the generation of surface fluxes (water, energy and CO₂) inside an ecological system are associated with the physical structure of the canopy and the amount of green biomass (biological component) which regulate both the radiation balance within the canopy and canopy-atmosphere energy and CO₂ exchanges. As such, one of the key parameters in the estimation and quantification of these fluxes is Leaf Area Index (LAI), defined as leaf area per unit of ground area (Montheith, 1973). The effect of perturbations such as logging on these processes is not well understood. This study examines the seasonal variability of litter biomass (leaves, flowers, fruits, and wood) and LAI before and after selective logging in order to better understand the effects of logging on forest structure. This study is done in conjunction with the LBA eddy-flux covariance study “Measuring the Effects of Logging on the CO₂ and Energy Exchange of a Primary Forest in the Tapajós National Forest”.

Methods

Leaf biomass and Leaf Area Index:

In order to evaluate the production of leaves and other litter components, we installed 30 1m² litterfall traps (Fig.1), (the number recommended by Newbould (1997)), which were arrayed at 25-m intervals along two east-west transects in an 18-ha block (300m N-S, 600m E-W) upwind of the eddy covariance tower. Litter was collected bi-weekly (Fig.2) beginning in September 2000. The material was separated into leaves, wood and other parts (mainly flowers and fruits), oven dried and weighed. Prior to oven drying, the subsample's leaf area was determined using a computer scanner (scanner HP Scanjet 6300C, output resolution 300) and image processing software (Rootedge). The relationship between weight and area measured was calculated for each subsample and used to derive an equation (Fig.4) which is used to estimate total leaf area for each litterfall trap.



Figure 2: Litterfall collection



Figure 1: Litterfall trap



Figure 3: Samples of leaves

Pre-Logging Results

Figure 5. Litterfall - FLONA Tapajós Km83




Figure 3. Area x weight





Figure 6. Fallen leaves x Rainfall - FLONA Tapajós km 83



Preliminary Post-Logging Results

Preliminary results after logging show a fall in leaf litter production of about 13,5% compared to the same period before selective logging (Fig.6).

This result corresponds with the results of hemispherical photography that found a decrease from 5 LAI to a mean of 4,3 m²m⁻² in the 3 months following the cut.

We will continue to monitor post-logging litterfall and LAI dynamics.

Acknowledgments

Antonio Oviedo for advice on collection methodology, Jose Mauro Moura for assistance with field work, CNPq and LBA for financial support.

References:
 Montheith, J.L. 1973. Principles of environmental physics. Edward Arnold ed. London
 Newbould, P.J., 1967. Methods of estimating the primary production of forests. IBP Handbook N° 2, Blackwell Scientific Publication, Oxford.
 Roberts, J.M. Cabral, G.R.M. da Costa, J.P.; McWilliam, A.L.C., Sa, T.D.A. 1996. An overview of the leaf area index and physiological measurements during ABRACOS. In: Gash, J.C.; Nobre, C.; Roberts J.M.; Vitoria R.L. (eds) 1996. Amazonian deforestation and climate. John&Wiley&Sons, UK. 61p.

Figure 7. Leaf fall before and after logging

