

Dependence of NAO variability on coupling with sea ice

Courtenay Strong · Gudrun Magnusdottir

Received: date / Accepted: date

Abstract The variance of the North Atlantic Oscillation index (denoted N) is shown to depend on its coupling with area-averaged sea ice concentration anomalies in and around the Barents Sea (index denoted B). The observed form of this coupling is a negative feedback whereby positive N tends to produce negative B , which in turn forces negative N . The effects of this feedback in the system are examined by modifying the feedback in two modeling frameworks: a statistical vector autoregressive model (F_{VAR}) and an atmospheric global climate model (F_{CAM}) customized so that sea ice anomalies on the lower boundary are stochastic with adjustable sensitivity to the model's evolving N . Experiments show that the variance of N decreases nearly linearly with the sensitivity of B to N , where the sensitivity is a measure of the negative feedback strength. Given that the sea ice concentration field has anomalies, the variance of N

Courtenay Strong

University of Utah, Salt Lake City, Utah, 84112, USA

E-mail: court.strong@utah.edu

Gudrun Magnusdottir

University of California, Irvine, Irvine, California, 92697-3100, USA

goes down as these anomalies become more sensitive to N . If the sea ice concentration anomalies are entirely absent, the variance of N is even smaller than the experiment with the most sensitive anomalies. The results have implications for simulated N variability in atmospheric global climate models, and also for the predictability of N under projected decreases in winter sea ice extent.

1 Introduction

Physical reasoning suggests that winter sea ice variability over the North Atlantic should be sensitive to the overlying atmospheric circulation since the latter can generate anomalies of sea ice velocity, atmospheric heat transport, and oceanic heat transport. As an example of this coupling, the upward trend of the North Atlantic Oscillation (NAO) index (N) from the 1960s through the mid 1990s increased the rate of winter sea ice retreat over the North Atlantic (Deser, 2000; Venegas and Mysak, 2000; Rigor et al, 2002; Hu et al, 2002; Liu and Curry, 2004; Rothrock and Zhang, 2005; Ukita et al, 2007). Since then, the NAO trend has reversed, and an overall downward trend in total sea ice extent is emerging that appears to be anthropogenic (Johannessen et al, 2004) and accelerating in summer (e.g. Comiso, 2006; Serreze et al, 2007). There nonetheless remains, superimposed on this overall downward trend in sea ice extent, a measurable signature of forcing by atmospheric circulation variability (Comiso, 2006; Maslanik et al, 2007; Francis and Hunter, 2004; Deser and Teng, 2008).

A substantial fraction of this atmospheric forcing is connected to the NAO, whose imprint is discernible as wind-driven sea ice extent anomalies in daily data (Kimura and Wakatsuchi, 2001), and ice motion and thickness anomalies in multi-year satellite records (Kwok et al, 2005). During positive NAO, sea ice concentrations in the Barents

Sea tend to be lower than average in association with increased temperatures related to enhanced atmospheric and oceanic heat transport (Yamamoto et al, 2006; Wang et al, 2000; Liu and Curry, 2004). Koenigk et al (2009) recently used a fully coupled model to show that that sea ice concentrations within the Barents Sea are most sensitive to wind-driven sea ice transport, with oceanic heat transport playing a small role in interannual variations and a large role in variations on longer time scales.

NAO-driven sea ice variations project strongly onto, and are likely largely responsible for, the leading pattern of North Atlantic sea ice concentration variability (Deser, 2000). This leading pattern consists of a dipole pattern of oppositely signed concentration anomalies in the Labrador and Barents Seas, where Barents Sea concentrations are lower during positive NAO. We refer to this variability pattern as the Greenland Sea-ice Dipole (GSD).

Modeling studies have shown that a positive GSD-like sea ice pattern sustained from December through April will generate a negative NAO-like hemispheric-scale response (Magnusdottir et al, 2004; Deser et al, 2004; Alexander et al, 2004; Kvamsto et al, 2004). These results evidence the presence of a negative feedback since the positive NAO produces a positive GSD pattern. Deser et al (2007) showed that this negative feedback begins as a baroclinic response localized to the forcing that reaches peak intensity in 5-10 days and persists for 2-3 weeks. If the GSD pattern is sustained beyond several weeks, the atmosphere develops a larger-scale equivalent barotropic response resembling the negative polarity of the Northern Annular Mode, which is maintained primarily by nonlinear transient fluxes of eddy vorticity (Deser et al, 2007) related in part to changes in Rossby wave breaking (Strong and Magnusdottir, 2009c).

There is evidence of non-stationarity in the association between sea ice and the NAO related to multi-decadal external forcing. Modeling studies show, for example,

a strong impact of the North Atlantic meridional overturning circulation (MOC) on sea ice in the Arctic Ocean and Barents Sea (Delworth et al, 1997; Jungcalus et al, 2005). Reconstructing North Atlantic sea ice extent back to 1800, Fauria et al (2009) found weak running correlations prior to 1950 and significant correlations thereafter. For the period 1978-2007, Strong and Magnúsdóttir (2009b) (hereafter, SM9) detected significant negative feedback between winter sea ice and the NAO at weekly time scales using satellite observations, reanalysis data, and the testable definitions of causality and feedback developed by Granger (1969). For interannual and longer time scales, Strong and Magnúsdóttir (2009a) examined multi-model ensemble simulations of the 20th to 23rd centuries, and found that an NAO-driven pattern of sea ice variability will persist but change somewhat in form as the ice edge retreats under projected global warming.

The present manuscript is focused on how the feedback between sea ice and the NAO affects the variance of sea ice and the variance of the NAO index. If, for example, we turn off the feedback or double the sensitivity of the feedback, how is the variance of the NAO index affected? To quantify the effects of the sea ice-NAO feedback, we use two observationally-motivated models of the sea ice-NAO system. The first model is based on the vector-autoregressive statistical model used in SM9. The second model is an atmospheric global climate model modified so that sea ice anomalies on the lower boundary are stochastic with adjustable sensitivity to the model's evolving NAO index. Our observational data are described in Section 2, followed by our modeling methods (Section 3), Results (Section 4), and a Summary and Discussion (Section 5).

2 Data

Consistent with SM9, we defined the NAO index N based on the leading empirical orthogonal function (EOF) of weekly mean NCEP/NCAR reanalysis sea level pressure data for the 21-week extended winter from 4 December through 23 April for years 1978-2008. Data were detrended, deseasonalized, and restricted to the domain used in Hurrell (1995) (20° - 80° N and 90° W- 40° E). A portion of this EOF is contoured in Fig. 1.

SM9 used a sea ice index (G) based on the GSD pattern. Here, we simplify this by focusing on the Barents Sea center of action since it accounts for nearly the entire feedback signal detected by Magnusdottir et al (2004). We define an index B which is the area-weighted, weekly-mean, sea ice concentration anomaly within the region outlined in Fig. 1, where the anomaly is relative to the long term mean for that week. This definition for B yields an intuitive sign convention whereby high B corresponds to anomalously high sea ice concentrations over the Barents Sea, but we note that the sign of B is opposite to the sign of the GSD index, so the negative feedback in this system is

$$\text{positive } N \rightarrow \text{negative } B \rightarrow \text{negative } N \quad (1)$$

as shown in Section (4.1).

The B index is calculated using National Snow and Ice Data Center (NSIDC) sea ice concentrations derived from Nimbus-7 Scanning Multichannel Microwave Radiometer and Defense Meteorological Satellite Program Special Sensor Microwave / Imager radiances (Cavalieri et al, 2008). These sea ice data are on a 25-km grid nominally once every two days for 1978–1986 and once daily for 1987 to present. We do not include winter 1987-1988 in our study because of a data gap.

3 Modeling

We use two modeling frameworks to explore how feedback between sea ice and the NAO affects the variance of sea ice and the variance of the NAO. The first framework is a linear statistical model coupling N and B as a vector-autoregressive (VAR) process (Section 3.1, framework denoted F_{VAR}). The second framework couples a linear stochastic model of B to the NCAR Community Atmosphere Model (CAM) Version 3.0 (Section 3.2, framework denoted F_{CAM}). The simplicity of F_{VAR} allows us to run many long experiments with minimal computational expense, and to obtain explicit expressions for how the variances of N and B are affected by feedback. The F_{VAR} results provide a view of system behavior based on a linear framework, and we compare these linear results to analogous experiments in the F_{CAM} system which is solving nonlinear partial differential equations to determine N .

3.1 Statistical model

In SM9, we studied feedback between sea ice and the NAO using a VAR model that included contemporaneous as well as lagged effects. For our purposes here, we maintain lag order p , simplify the model by excluding contemporaneous effects, and introduce “feedback scaling parameters” g and h to be used when experimenting with the model.

Denoting N during week t by N_t , and B during week t by B_t , we write F_{VAR} as

$$\begin{pmatrix} B_t \\ N_t \end{pmatrix} = \sum_{i=1}^p \begin{pmatrix} \alpha_i & g\beta_i \\ h\eta_i & \phi_i \end{pmatrix} \begin{pmatrix} B_{t-i} \\ N_{t-i} \end{pmatrix} + \begin{pmatrix} \varepsilon_{Bt} \\ \varepsilon_{Nt} \end{pmatrix} \quad (2)$$

where B_t and N_t are stationary with zero mean, and ε_{Bt} and ε_{Nt} are uncorrelated white noise disturbances with respective standard deviations σ_B and σ_N . The parameters g and h govern, respectively, how sensitive B is to N and how sensitive N is to B . In

general, and particularly when fitting F_{VAR} to observations, $g = h = 1$. The feedback scaling parameters may be given values other than 1 for the purpose of experimentation aimed at determining the response of the system to a stronger (e.g., $g = 1.5$) or weaker feedback (e.g., $g = 0.5$). We can write (2) compactly as

$$\mathbf{x}_t = \sum_i \Phi_i \mathbf{x}_{t-i} + \varepsilon_t. \quad (3)$$

To determine the appropriate order p , we fit the model to observations at order p (i.e., “unrestricted”) and compare it to the model fit at order $p - 1$ (i.e., “restricted”). Where we detect a significant degradation in model strength going from p to $p - 1$, we declare p to be the appropriate model order. Significant degradations in model strength are tested for using the log-likelihood ratio given by Sims (1980)

$$L \equiv (T - c)(\log |\boldsymbol{\Sigma}_r| - \log |\boldsymbol{\Sigma}_u|) \quad (4)$$

where T is the number of use-able observations, c is the maximum number of regressors in the longest equation, and $|\boldsymbol{\Sigma}_u|$ and $|\boldsymbol{\Sigma}_r|$ are the determinants of the covariance matrices of the unrestricted and restricted model’s residuals, respectively.

To quantify the variance of N and the variance of B in F_{VAR} , we want expressions for the $2p + 1$ covariance matrices

$$\boldsymbol{\Gamma}(m) \equiv \text{E}(\mathbf{x}_{t+m} \mathbf{x}_t^T) = \begin{pmatrix} \gamma_{BB}(m) & \gamma_{BN}(m) \\ \gamma_{NB}(m) & \gamma_{NN}(m) \end{pmatrix}, \quad m = 0, \pm 1, \dots, \pm p \quad (5)$$

where m is lag, $(\cdot)^T$ denotes transpose, $\text{E}(\cdot)$ denotes expectation, and $\gamma_{BN}(m)$ is the covariance where N leads B by m weeks. For convenience, we denote the variances of N and B as γ_{NN} and γ_{BB} . The $2p + 1$ matrices in (5) have a total of $8p + 4$ elements, but at most $4p + 3$ of them are unique since $\boldsymbol{\Gamma}(-m) = \boldsymbol{\Gamma}^T(m)$ and $\gamma_{NB}(0) = \gamma_{BN}(0)$. To get $4p + 3$ linear equations for these $4p + 3$ unique matrix elements, we post-multiply

(3) by $\mathbf{x}_{t-j}^T, j = 0, 1, \dots, p$ and take expectations, yielding (e.g., Brockwell and Davis, 1996)

$$\boldsymbol{\Gamma}(i) = \sum_{j=1}^p \boldsymbol{\Phi}_j \boldsymbol{\Gamma}(i-j), \quad i = 1, 2, \dots, p \quad (6)$$

$$\boldsymbol{\Gamma}(0) = \sum_{j=1}^p \boldsymbol{\Phi}_j \boldsymbol{\Gamma}(-j) + \boldsymbol{\Upsilon} \quad (7)$$

where $\boldsymbol{\Upsilon}$ is the covariance matrix of ε_t . Numerically solving the linear system (6-7) yields all the unique elements of (5), and we will be focusing primarily on the variances γ_{NN} and γ_{BB} .

3.2 Atmospheric global climate model

As described above, F_{VAR} couples a linear expression for N to a linear expression for B where the latter is written

$$B_t = \sum_{i=1}^p (\alpha_i B_{t-i} + g\beta_i N_{t-i}) + \varepsilon_{Bt}. \quad (8)$$

F_{CAM} retains (8) as the linear expression for B , but couples it to CAM. To achieve this coupling, we wrote a parallelized module for CAM that introduces a weekly sea ice concentration anomaly according to equation (8), but with the values of weekly mean N calculated from the sea level pressure fields being generated within CAM during the run. The module requires three input files: 1) the spatial pattern of the NAO obtained from a long unforced run analyzed the same way as the observational NAO (Section 2), 2) a weekly climatology of sea level pressure based on a long unforced run, and 3) a “ B -file” containing sea ice concentration anomaly fields for a range of B values and months.

The sea ice concentration anomaly fields in the B -file are specified as a function of time, latitude, longitude, and the index B . Each field is a composite of NSIDC sea

ice concentration anomaly observations grouped by the five months December through April, and seven B index bins centered around the integers $-3, -2, \dots, 3$. As illustrative examples, Fig. 1a shows the composite sea ice anomaly for all January observations with B indices in the bin centered on $B = -2$, and Fig. 1b shows the composite sea ice anomaly for all February observations with B indices in the bin centered on $B = 3$. The B -file anomalies are observations interpolated onto the model grid. Experimentation with F_{CAM} using more idealized, smooth anomalies produced results similar to those presented here.

Week $t = 1$ is defined as the first seven days of model integration, and F_{CAM} specifies the preceding p weeks as initial conditions. When the model initializes, it therefore sets $t = 1$ and does the following:

1. Set initial conditions $B_{t=1-i} = 0$ for $i = 1, 2, \dots, p$.
2. Set initial conditions $N_{t=1-i} = c$ for $i = 1, 2, \dots, p$ where c is the value of the NAO index calculated from the sea level pressure field in the initial condition file.
3. Calculate B_1 using the $2p$ initial conditions and equation (8).
4. Determine the sea ice concentration anomaly field to be applied during week $t = 1$ by going to the current model month in the B -file and linearly interpolating the sea ice concentration anomaly at each grid point as a function of B to the value B_1 .

The module then takes the steps required to perform the following every seven days beginning on week $t = 2$:

1. Calculate N_{t-1} using the preceding week's sea level pressure fields, which the model has stored, weighted by the square root of the grid area, deseasonalized, and projected onto the spatial pattern of the NAO.

-
2. Calculate B_t from equation (8) using B_{t-i} and N_{t-i} for $i = 1, 2, \dots, p$.
 3. Determine the sea ice concentration anomaly field to be applied during next seven days by going to the current model month in the B -file and linearly interpolating the sea ice concentration anomaly at each grid point as a function of B to the value B_t .

For the sea ice concentration anomalies used in our experiments, going to a specific month in the B -file produces anomaly patterns that are similar to those obtained by performing a more expensive bi-linear interpolation as a function of month and B_t .

F_{CAM} can be thought of as a modeling framework that is intermediate between running CAM coupled to a full ice model and running CAM forced by sea ice linearly interpolated from a climatology. This intermediate framework is useful for our purposes because we can explicitly control aspects of B including whether, and to what degree, B is sensitive to variations in CAM’s evolving N .

3.3 Experiments

We designed our experiments to uncover the effects of feedback in the B and N system. In the experiments, we varied the values of the feedback scaling parameters g and h as shown in Table 1. Our control case (CTL) corresponds conceptually to the observed system, with N and B having realistic sensitivity to one another (i.e., $g = h = 1$). For the CTL case, B evolved as a vector autoregressive process sensitive to the past states of itself, the past states of N , and a stochastic forcing. For the IND experiment, $g = h = 0$ meaning that B and N evolved independently. In terms of F_{CAM} , the IND case is equivalent to forcing the atmosphere with an anomaly-free sea ice climatology. For the AR experiment, B evolved as an autoregressive process, meaning it was sensitive

to the past states of itself and a stochastic forcing. For the VAR2 experiment, B evolved as in CTL but with a doubled sensitivity to N (i.e., $g = 2$). In the physical system, a change in the responsiveness of B to N could be related to, for example, changes in the thickness or extent of the perennial ice.

For F_{CAM} , we developed a 150-member ensemble for each of CTL, IND, AR, and VAR2. Each member covered the 21 weeks beginning on December 4, with initial conditions taken from a long unforced run. We define the response as the total variance of the experiment ensemble divided by the total variance of the CTL ensemble. For F_{VAR} , we define the response as the variance in the experiment case divided by the variance of the CTL case, where each variance comes from the solution to the linear system (6-7). We denote the response by a vertical bar followed by a subscript denoting the experiment. For example, the response of γ_{NN} in the AR experiment is $\gamma_{NN}|_{\text{AR}}$.

4 Results

We have two results sections. In the first (4.1), we present observations of N and B , fit the VAR to these observations, and verify that the F_{VAR} and F_{CAM} models reasonably capture the observed behavior of B and N . In Section 4.2, we present results from experimentation with F_{VAR} and F_{CAM} .

4.1 Observations

For observations, the lagged correlations of N and B are shown in Fig. 2, and we will interpret them in the context of a hypothetical, anomalously high value of N . Concurrent with this high value of N (at lag 0), B tends to be anomalously low, indicating a reduction in sea ice over the Barents Sea that is physically consistent

with the patterns of temperature advection and sea ice velocity associated with the positive NAO. This tendency for anomalously low B is visible over lags of several weeks forward from the N anomaly. One to six weeks after B is anomalously low, N tends to decrease (positive correlations toward left side of Fig. 2). This is evidence of the negative feedback between N and B discussed in the Introduction. Comparison of Figs. 2b,c shows that B is more autocorrelated than N and subject to less short-term fluctuations or “noise.”

Fitting system (2) to observations, we find the appropriate model order to be $p = 4$ (method in Section 3.1) as in SM9. Parameter values for Φ matrix are as given in Table 2. By testing the significance of these parameters using equation 4, we conclude at the 95% confidence level that there is Granger feedback (Granger, 1969) between N and B . SM9 provide a closely related result and more discussion of Granger feedback detection in this application. The blue and red curves in Fig. 2 show, respectively, lagged correlations for output from F_{VAR} and F_{CAM} . Both models capture the temporal covariation of N and B reasonably well.

4.2 Model responses

We first use F_{VAR} to show how γ_{NN} and γ_{BB} respond to the g and h feedback scaling parameter settings in the IND, AR, and VAR2 experiments. As noted in Section 3.1, we can directly obtain these responses by solving the linear system (6-7) with different values for g and h . For the CTL case, we solved equations (6-7) with $g = h = 1$, and Table 3 provides some select values from this solution: γ_{NN} , γ_{BB} , and γ_{BN} (3). Fig. 3 shows that synthetic data generated from F_{VAR} converge toward these numerical solutions as t becomes large.

To provide a context for the three experiment results, we calculated γ_{NN} and γ_{BB} responses for a large set of values of g and h ranging from -1 to 2.5 (Figs. 4a and 4b, respectively). Fig. 4c is a schematic indicating key locations or regions in the hg response plane. The open circles mark the locations of the CTL case and the three experiments IND, AR, and VAR2. The responses in Figs. 4a and 4b are both 1.0 for the CTL case by definition.

In the AR experiment, feedback is turned off by setting $g = 0$ meaning that B is independent of previous values of N , but N still depends on past values of B . The response is a 5% increase in γ_{NN} (i.e., $\gamma_{NN}|_{AR} = 1.05$, Fig. 4a). To provide the details underlying $\gamma_{NN}|_{AR}$, we write the equation for γ_{NN} from (7)

$$\gamma_{NN} = \sum_{i=1}^4 [h\eta_i\gamma_{NB}(i) + \phi_i\gamma_{NN}(i)] + \sigma_N^2. \quad (9)$$

The $\gamma_{NB}(i)$ terms all become positive in AR meaning that anomalies of B tend to be followed by like-signed anomalies of N , and the $\gamma_{NN}(i)$ terms increase, meaning that the autocorrelation of N increases. Both changes contribute toward higher γ_{NN} . In the same experiment, the variance of B decreases by 6% ($\gamma_{BB}|_{AR} = 0.94$, Fig. 4b). In the equation for γ_{BB} from (7)

$$\gamma_{BB} = \sum_{i=1}^4 [\alpha_i\gamma_{BB}(i) + g\beta_i\gamma_{BN}(i)] + \sigma_B^2, \quad (10)$$

setting $h = 0$ reduces the variance of B because the γ_{BN} terms contribute positively to γ_{BB} in CTL.

In the VAR2 experiment, we increase the strength of the negative feedback by setting $g = 2$. The variance, γ_{NN} , decreases by approximately 2% ($\gamma_{NN}|_{VAR2}=0.98$, Fig. 4a) and γ_{BB} increases by approximately 42% ($\gamma_{BB}|_{VAR2}=1.42$, Fig. 4b). In the IND experiment, the settings $g = h = 0$ isolate N and B from each other, and the variance of each decreases ($\gamma_{NN}|_{IND}=0.99$ in Fig. 4a, $\gamma_{BB}|_{IND}=0.93$ in Fig. 4b).

Commenting more generally on the surfaces in Fig. 4, the responses become rapidly large in portions of the gh plane that are shaded in Fig. 4c. In these shaded regions, the sign of either g or h becomes negative, rendering the feedback positive and generating very large variances of N and B . The response surfaces are symmetrical about the origin with asymptotes (dashed lines) along which the contoured response is isolated from the other system variable, as in the IND case. Using the γ_{NN} response as an example (Fig. 4a), there is an asymptote at $h=0$ because this scaling desensitizes N to B , so a change in g , which is a scaling factor in the B equation, has no effect on N .

We now turn to the F_{CAM} results (red circles, Fig. 5). For the purposes of comparison, Fig. 5 shows portions of the F_{VAR} results as curves. These F_{VAR} response curves are taken from the surfaces in Fig. 4 running down the plots from VAR2 to CTL to AR (curve in Fig. 5a taken from Fig. 4a; curve in Fig. 5b taken from Fig. 4b). Considering the AR, CTL, and VAR2 results, the F_{CAM} results agreed qualitatively with the F_{VAR} results: turning feedback off in AR increases γ_{NN} and decreases γ_{BB} , whereas doubling the feedback sensitivity in VAR2 produces the opposite responses. The dashed lines in Fig. 5 show that γ_{NN} responded nearly linearly to g ($r^2 > 0.99$, slope -0.04), as did γ_{BB} ($r^2 > 0.95$, slope 0.32). We conclude that the variance of N and B are approximately linear functions of the sensitivity of B to N for the range $0 \leq g \leq 2$.

For the IND case, the γ_{BB} response is the same as in the AR case (not shown) because B lacks sensitivity to N in both experiments (i.e., $g = 0$, Table 1). For γ_{NN} in the IND case, the F_{CAM} result matches in sign, but is stronger than the F_{VAR} result (Fig. 5a). Specifically, F_{VAR} predicts a small γ_{NN} response when isolating N from B , but F_{CAM} produces the strongest response in γ_{NN} for the cases we examined, amounting to a 6% decrease. F_{VAR} and F_{CAM} are thus in agreement about responses with respect to the sensitivity of B to N (i.e., g), but agree less well about responses with respect to

the sensitivity of N to B (i.e., h). This is not entirely unexpected since F_{CAM} generates N using nonlinear equations of motion and parameterized model physics.

5 Summary and discussion

We quantified the effects of negative feedback between the North Atlantic Oscillation index (denoted N) and an index of sea ice concentration anomalies in and around the Barents Sea (denoted B). Statistically testable definitions of causality and feedback were used to conclude that, in observations, positive N tends to produce negative B , which in turn forces negative N . We then investigated this feedback by modifying it in two modeling frameworks: a statistical vector autoregressive model (F_{VAR}) and an atmospheric global climate model (F_{CAM}) customized so that sea ice anomalies on the lower boundary were stochastic with adjustable sensitivity to the model’s evolving N . We focused on how the variance of N and the variance of B respond to changes in the system feedback.

We defined a control case (CTL) in which B evolved as a vector autoregressive process sensitive to the past states of itself, the past states of N , and a stochastic forcing. For the IND experiment, B and N evolved independently. For the AR experiment, B evolved as an autoregressive process, meaning it was sensitive to the past states of itself and a stochastic forcing. For the VAR2 experiment, B evolved as in CTL but with a doubled sensitivity to N . In the physical system, a change in the responsiveness of B to N could be related to, for example, changes in the thickness or extent of the perennial ice.

In the following conclusions, we take “feedback strength” to mean the value of the feedback scaling parameter g , which is the sensitivity of B to N . The variance of N

(γ_{NN}) scales approximately linearly with a negative slope ($r^2 > 0.99$, slope -0.04) with feedback strength in F_{CAM} , in reasonable agreement with F_{VAR} results. The variance of B (γ_{BB}) scales approximately linearly ($r^2 > 0.95$, slope 0.32) with feedback strength, also in reasonable agreement with F_{VAR} results. In F_{CAM} , the IND case produced the strongest response in γ_{NN} , amounting to a 6% decrease in variance. This is different from the F_{VAR} prediction that γ_{NN} would be very similar in the IND and CTL cases. This difference indicates that F_{VAR} and F_{CAM} produce reasonably similar responses when feedback is scaled by changing the sensitivity of B to N (i.e., g), but produce less similar responses when feedback is scaled by changing the sensitivity of N to B .

Based on the F_{CAM} results, the variance of N increased progressively from IND to VAR2 to CTL to AR. In other words, a zero-anomaly sea ice climatology (IND case) produces minimal N variance whereas, given that the sea ice field has anomalies (VAR2, CTL, or AR cases), the variance of N goes down as these anomalies become more sensitive to N . This negative-slope linear response of γ_{NN} to the feedback strength is consistent with the fundamental behavior of negative feedback, and its quantification has implications for the simulation of internal variability in atmospheric global climate models forced by sea ice, and for the predictability of N under projected changes in winter sea ice extent.

Acknowledgements This research was supported by NSF grant ATM-0612779 and NOAA grants NA06OAR4310149 and NA09OAR4310132.

References

Alexander MA, Bhatt US, Walsh JE, Timlin MS, Miller JS, Scott JD (2004) The atmospheric response to realistic Arctic sea ice anomalies in an AGCM during winter.

-
- J Climate 17:890–905
- Brockwell PJ, Davis RA (1996) Introduction to Time Series and Forecasting. Springer-Verlag, New York
- Cavalieri D, Parkinson C, Gloersen P, Zwally HJ (2008) Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I passive microwave data, 1978-2008. National Snow and Ice Data Center, Boulder, Colorado. Digital media
- Comiso JC (2006) Abrupt decline in arctic winter sea ice cover. Geophys Res Lett 33, DOI 10.1029/2006GL027341
- Delworth TL, Manabe S, Stouffer RJ (1997) Multidecadal climate variability in the Greenland Sea and surrounding regions: a coupled model study. Geophys Res Lett 24:257–260, DOI 10.1029/96GL03927
- Deser C (2000) On the teleconnectivity of the “Arctic Oscillation”. Geophys Res Lett 27:779–782
- Deser C, Teng H (2008) Evolution of Arctic sea ice concentration trends and the role of atmospheric circulation forcing, 1979-2007. Geophys Res Lett 35, DOI 10.1029/2007GL032023
- Deser C, Magnusdottir G, Saravanan R, Phillips AS (2004) The effects of North Atlantic SST and sea-ice anomalies on the winter circulation in CCM3. Part II: Direct and indirect components of the response. J Climate 17:877–889
- Deser C, Tomas RA, Peng S (2007) The transient atmospheric circulation response to North Atlantic SST and sea ice anomalies. J Climate 20:4751–4767
- Fauria MM, Grinsted A, Helama S, Moore J, Timonen M, Martma T, Isaksson E, Eronen E (2009) Unprecedented low twentieth century winter sea ice extent in the Western Nordic Seas since A.D. 1200. Clim Dyn DOI 10.1007/s00382-009-0610-z

- Francis JA, Hunter E (2004) Drivers of declining sea ice extent in the Arctic winter: a tale of two seas. *Geophys Res Lett* 31:145–160
- Granger CWJ (1969) Investigating causal relationships by econometric models and cross-spectral methods. *Econometrica* 37:473–485
- Hu A, Rooth C, Bleck R, Deser C (2002) Nao influence on sea ice extent in the eurasian coastal region. *Geophys Res Lett* 29, DOI 10.1029/2001GL014293
- Hurrell JW (1995) Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation. *Science* 269:676–679
- Johannessen OM, Bengtsson L, Miles MW, Kuzmina SI, Semenov VA, Alekseev GV, Nagurnyi AP, Zakharov VF, Boblyev LP, Petterson LH, Hasselmann K, Cattle HP (2004) Arctic climate change: observed and modelled temperature and sea-ice variability. *Tellus* 56:328–341
- Jungclauss JH, Haak H, Latif M, Mikolajewicz U (2005) Arctic-North Atlantic interactions and multidecadal variability of the Meridional Overturning Circulation. *J Climate* 18:4013–4031
- Kimura N, Wakatsucchi M (2001) Mechanism for the variation of sea ice extent in the Northern Hemisphere. *J Geo Res* 106:31,319–31,331
- Koenigk T, Mikolajewicz U, Jungclauss JH, Kroll A (2009) Sea ice in the Barents Sea: seasonal to interannual variability and climate feedbacks in a global coupled model. *Clim Dyn* 32:1119–1138
- Kvamsto NG, Skeie P, Stephenson DB (2004) Impact of Labrador sea-ice extent on the North Atlantic Oscillation. *Int J Climatol* 24:603–612
- Kwok R, Maslowski W, Laxon SW (2005) On large outflows of Arctic sea ice into the Barents Sea. *Geophys Res Lett* 32, DOI 10.1029/2005GL024485

-
- Liu J, Curry JA (2004) Recent Arctic sea ice variability: connections to the Arctic Oscillation and the ENSO. *Geophys Res Lett* 31, DOI 10.1029/2004GL019858
- Magnusdottir G, Deser C, Saravanan R (2004) The effects of North Atlantic SST and sea ice anomalies on the winter circulation in CCM3. Part I: main features and storm track characteristics of the response. *J Climate* 17:857–876
- Maslanik J, Drobot S, Fowler C, Emery W, Barry R (2007) On the Arctic climate paradox and the continuing role of atmospheric circulation in affecting sea ice conditions. *Geophys Res Lett* 34, DOI 10.1029/2006GL028269
- Rigor IG, Wallace JM, Colony RL (2002) Response of sea ice to the arctic oscillation. *J Climate* 15:2648–2663
- Rothrock DA, Zhang J (2005) Arctic ocean sea ice volume: What explains its recent depletion? *J Geo Res* 110, DOI 10.1029/2004JC002282
- Serreze MC, Holland MM, Stroeve J (2007) Perspectives on the arctic’s shrinking sea-ice cover. *Science* 315:1533–1536
- Sims CA (1980) Macroeconomics and reality. *Econometrica* 48:1–48
- Strong C, Magnusdottir G (2009a) Modeled winter sea ice variability and the North Atlantic Oscillation: a multi-century perspective. *Clim Dyn* DOI 10.1007/s00382-009-0550-7
- Strong C, Magnusdottir G (2009b) Observed feedback between winter sea ice and the North Atlantic Oscillation. *J Climate* In press
- Strong C, Magnusdottir G (2009c) The role of Rossby wave breaking in the Pacific Decadal Oscillation. *J Climate* 22:1819–1833
- Ukita J, Honda M, Nakamura H, Tachibana Y, Cavalieri DJ, Parkinson CL, Koide H, Yamamoto K (2007) Northern Hemisphere sea ice variability: Lag structure and its implications. *Tellus, Ser A* 59:261–272, DOI 10.1111/j.1600-0870.2006.0223.x

Venegas SA, Mysak LA (2000) Is there a dominant timescale of natural climate variability in the arctic? *J Climate* 13:3412–3434

Wang J, Wu B, Tang C, Walsh JE, Ikeda M (2000) Seesaw structure of subsurface temperature anomalies between the Barents Sea and the Labrador Sea. *Geophys Res Lett* 31, DOI 10.1029/2004GL019981

Yamamoto K, Tachibana Y, Honda M, Ukita J (2006) Possible feedback of winter sea ice in the Greenland and Barents Seas on the local atmosphere. *Geophys Res Lett* 33, DOI 10.1029/2006GL026286




Fig. 1 Shading shows example sea ice anomaly fields from the “B-file” valid for (a) January with $-3.5 \leq B < -2.5$ and (b) February with $2.5 \leq B < 3.5$. The EOF of the NAO is contoured with negative values dashed and the zero contour suppressed.




Fig. 2 Lagged Pearson correlation coefficients r for (a) B and N , (b) N with itself, and (c) B with itself. In each panel, the black curve with filled circles represents observations, the blue curve represents F_{VAR} model output, and the red curve represents F_{CAM} model output.




Fig. 3 Output from F_{VAR} converging to solutions of linear system (6-7) as t becomes large. Variables are (a) the variance of N , (b) the variance of B , and (c) the covariance where N leads B by three weeks. The red line shows the solutions and the black curve shows the mean of 100 simulations. The distribution of results at each week t is shaded in 10-percentile increments (i.e., yellow is 10-90th percentile, light green is 20-70th percentile, etc.). The vertical lines at $t = 3150$ show the number of weeks in each F_{CAM} ensemble.




Fig. 4 The response of (a) the variance of N (denoted γ_{NN}) and (b) the variance of B (denoted γ_{BB}) to changes in the feedback scaling parameters g and h , and (c) a schematic indicating key areas in the hg plane. Further details in the text.




Fig. 5 (a) For the variance of N (denoted γ_{NN}), the curve and cross show the responses given by F_{VAR} for the four cases indicated on the abscissa. The red circles show the responses produced by F_{CAM} with error bars showing plus or minus one standard error. The dashed line is a least squares linear fit to the AR, CTL and VAR2 results from F_{CAM} as a function of g . (b) Same as (a), but for the variance of B (denoted γ_{BB}), and the IND case is not shown because it is equivalent to the AR case.

Table 1 For each experiment: a description, the settings of the scaling parameters g and h (see equation 2), and the notation used to refer to the experiment.

Description	g	h	notation
independent B and N	0	0	IND
autoregressive B	0	1	AR
vector autoregressive B	1	1	CTL
enhanced vector autoregressive B	2	1	VAR2

Table 2 Values of the parameters in the Φ matrices of system (3) estimated from observations with $g = h = 1$. For the Υ matrix, the variance of ε_B is 0.40 and the variance of ε_N is 0.84.

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
α_i	0.95	-0.15	0.05	0.05
β_i	-0.16	0.06	-0.03	0.01
η_i	0.17	-0.21	0.31	-0.17
ϕ_i	0.55	-0.12	0.12	0.07

Table 3 Covariances for the system (3) solved for using observational estimates of the Φ and Υ matrices in Table 2.

γ_{NN}	1.02
γ_{BB}	0.84
$\gamma_{BN}(3)$	-0.18