

## Lecture 12: Future Climate Changes

- ❑ Intergovernmental Panel on Climate Change
- ❑ Main Conclusions of the SPM (Summary For Policy Makers)




## IPCC Web Site


<http://www.ipcc.ch>



## Structure of IPCC 1997 – 2001





(adapted from IPCC website; presented by John Houghton)



## History of IPCC

- 1988 - The United Nations Environment Programme (UNEP) and the World Meteorological Organization (WMO) establish the Intergovernmental Panel on Climate Change.
  - The United Nations General Assembly endorses the action of UNEP and the WMO in setting up the IPCC.
- 1990 - The IPCC publishes its First Assessment Report (Working Group I – *Climate Change: The IPCC Scientific Assessment*; Working Group II – *Climate Change: The IPCC Impacts Assessment*; Working Group III – *Climate Change: The IPCC Response Strategies*).
  - The UN General Assembly notes the report findings and decides to initiate negotiations for a framework convention on climate change.
- 1995 - The IPCC publishes its Second Assessment Report (Working Group I – *Climate Change 1995: The Science of Climate Change*; Working Group II – *Climate Change 1995: Impacts, Adaptations and Mitigation of Climate Change: Scientific-Technical Analyses*; Working Group III – *Climate Change 1995: Economic and Social Dimensions of Climate Change*; IPCC Second Assessment: *Climate Change 1995* (includes Synthesis Report)).
- 2001 - The IPCC publishes its Third Assessment Report (Working Group I – *Climate Change 2001: The Scientific Basis*; Working Group II – *Climate Change 2001: Impacts, Adaptation, and Vulnerability*; Working Group III – *Climate Change 2001: Mitigation*; *Climate Change 2001: Synthesis Report*).
- 2007 - The IPCC publishes its Fourth Assessment Report (ARA) (Working Group I – *Climate Change 2007: The Physical Science Basis*; Working Group II – *Climate Change 2007: Impacts, Adaptation and Vulnerability*; Working Group III – *Climate Change 2007: Mitigation of Climate Change*; *Climate Change 2007: Synthesis Report*).
- 2013 - The IPCC approves *Climate Change 2013: The Physical Science Basis*, the Working Group I contribution to AR5.





## IPCC AR5 (Fifth Assessment Report)

**WGI**  
(Sciences)

**WGII**  
(Impacts and Adaptation)

**WGIII**  
(Mitigation)

**Synthesis Report**

ESS15  
Prof. Jin-Yi Yu

## Climate Change 2001: The Scientific Basis

### WGI contribution to IPCC Third Assessment Report

**Summary for Policymakers (SPM)**

Approved 'sentence by sentence'  
by WGI plenary  
(99 Governments and 45 scientists)

**The Scientific Basis (Technical Summary)**

14 chapters; 120 Lead Authors  
515 Contributing Authors  
4621 References quoted

ESS15  
Prof. Jin-Yi Yu

### Contents

Foreword	vii
Preface	ix
Summary for Policymakers	1
Technical Summary	21
1 The Climate System: an Overview	85
2 Observed Climate Variability and Change	99
3 The Carbon Cycle and Atmospheric Carbon Dioxide	183
4 Atmospheric Chemistry and Greenhouse Gases	239
5 Aerosols, their Direct and Indirect Effects	289
6 Radiative Forcing of Climate Change	349
7 Physical Climate Processes and Feedbacks	417
8 Model Evaluation	471
9 Projections of Future Climate Change	525
10 Regional Climate Information – Evaluation and Projections	583
11 Changes in Sea Level	639
12 Detection of Climate Change and Attribution of Causes	695
13 Climate Scenario Development	739
14 Advancing Our Understanding	769
Appendix I Glossary	787
Appendix II SRES Tables	799
Appendix III Contributors to the IPCC WGI Third Assessment Report	827
Appendix IV Reviewers of the IPCC WGI Third Assessment Report	845
Appendix V Acronyms and Abbreviations	861
Appendix VI Units	869
Appendix VII Some Chemical Symbols used in this Report	871
Appendix VIII Index	873


ESS15  
Prof. Jin-Yi Yu

The IPCC Working Group \_\_\_\_\_ aims at assessing the physical scientific basis of the climate system and climate change.

- (A) I
- (B) II
- (C) III



## Understand and Project Our Climate System



## Major Conclusions in SPM (TAR)

1. An increasing body of observations gives a collective picture of a warming world and other changes in the climate system.
2. Emissions of greenhouse gases and aerosols due to human activities continue to alter the atmosphere in ways that are expected to affect the climate.
3. Confidence in the ability of models to project future climate has increased.
4. There is new and stronger evidence that most of the warming observed over the last 50 years is attributable to human activities.
5. Human influences will continue to change atmospheric composition throughout the 21st century.
6. Global average temperature and sea level are projected to rise under all IPCC SRES scenarios.
7. Anthropogenic climate change will persist for many centuries.
8. Further action is required to address remaining gaps in information and understanding.



## IPCC's Probability Phrases

Table 1 | IPCC guidelines for translation of probability phrases.

Phrase	Likelihood
Virtually certain	>99%
Very likely	>90%
Likely	>66%
About as likely as not	33%-66%
Unlikely	<33%
Very unlikely	<10%
Exceptionally unlikely	<1%

Note: In some IPCC reports authors have used the phrases 'More likely than not' for probabilities > 50%, 'Extremely likely' for probabilities above 95% and 'Extremely unlikely' for probabilities below 5%.



## How Future CO2 Emission Calculated?


$$\% \text{ increase in carbon emissions} = \% \text{ increase in population} \times \% \text{ change in emissions per person} \times \text{Changes in efficiency of carbon use}$$

- ❑ **Global Population:** is expected to increase to 11 billion between 2075 and 2100 (100% increase).
- ❑ **Emission Per Person:** is linked to averaged standard of living (such as car and home heating or cooling).
- ❑ **Efficiency of Use:** The hardest factor to project. The efficiency depends on technologies. This is also the factor that may keep carbon emissions from increasing in the future.





ESS15  
Prof. Jin-Yi Yu

## IPCC Special Report on Emission Scenarios (SRES)



ESS15  
Prof. Jin-Yi Yu


## IPCC Special Report on Emission Scenarios (SRES)



ESS15  
Prof. Jin-Yi Yu

## RCP vs. SRES

A "Parallel Approach" Implies Much More Interaction Between the IAV, IAM and CM communities



Prof. Jin-Yi Yu

## Representative Concentration Pathway (RCP)

Name	Radiative forcing	CO <sub>2</sub> equiv (p.p.m.)	Temp anomaly (°C)	Pathway	SRES temp anomaly equiv
RCP8.5	8.5 Wm <sup>2</sup> in 2100	1370	4.9	Rising	SRES A1F1
RCP6.0	6 Wm <sup>2</sup> post 2100	850	3.0	Stabilization without overshoot	SRES B2
RCP4.5	4.5 Wm <sup>2</sup> post 2100	650	2.4	Stabilization without overshoot	SRES B1
RCP2.6 (RCP3PD)	3Wm <sup>2</sup> before 2100, declining to 2.6 Wm <sup>2</sup> by 2100	490	1.5	Peak and decline	None

**Table 4:** from Moss et al. 2010. Median temperature anomaly over pre-industrial levels and SRES comparisons based on nearest temperature anomaly, from Rogelj et al. 2012


- RCPs are four greenhouse gas concentration (not emissions) trajectories adopted by the IPCC for its AR5 in 2014.
- It supersedes Special Report on Emissions Scenarios (SRES) projections published in 2000.
- RCP 4.5 is a scenario that stabilizes radiative forcing at 4.5 Watts per meter squared in the year 2100 without ever exceeding that value.




ESS15  
Prof. Jin-Yi Yu

## RCP vs. SRES

### Emission Scenarios




### Concentration Scenarios



ESS15  
Prof. Jin-Yi Yu

## Major Conclusions in SPM (TAR)

1. An increasing body of observations gives a collective picture of a warming world and other changes in the climate system.
2. Emissions of greenhouse gases and aerosols due to human activities continue to alter the atmosphere in ways that are expected to affect the climate.
3. Confidence in the ability of models to project future climate has increased.
4. There is new and stronger evidence that most of the warming observed over the last 50 years is attributable to human activities.
5. Human influences will continue to change atmospheric composition throughout the 21st century.
6. Global average temperature and sea level are projected to rise under all IPCC SRES scenarios.
7. Anthropogenic climate change will persist for many centuries.
8. Further action is required to address remaining gaps in information and understanding.



ESS15  
Prof. Jin-Yi Yu


## Evidences of Global Warming and Other Changes

- Temperature
- Precipitation
- Snow / Ice Cover
- Sea Level
- Circulation
- Extremes




ESS15  
Prof. Jin-Yi Yu

## Global Surface Temperature (TAR)




The global average surface temperature has increased over the 20th century by about 0.6°C.





ESS15  
Prof. Jin-Yi Yu

## Global Surface Temperature (AR5)




## Compared to the Past 1000 Years




ESS15  
Prof. Jin-Yi Yu

## The Land and Oceans Have Warmed



## The Land Have Warmed (AR5)


Observed change in surface temperature  
1901–2012



ESS15  
Prof. Jin-Yi Yu


## Oceans Have Warmed (AR5)

Observed change in surface temperature  
1901–2012



**North Atlantic  
“Warming Hole”**


→ may be due to the  
slow down of the  
AMOC.



- Ocean warming dominates the increase in energy stored in the climate system, accounting for more than 90% of the energy accumulated between 1971 and 2010 (high confidence).
- More than 60% of the net energy increase in the climate system is stored in the upper ocean (0–700 m) during the relatively well-sampled 40-year period from 1971 to 2010, and about 30% is stored in the ocean below 700 m.
- It is virtually certain that the upper ocean (0–700 m) warmed from 1971 to 2010.

## Precipitation Patterns Have Changed

Annual precipitation trends: 1900 to 2000




Rainfall maybe  
decreased over  
much subtropical  
lands.

**It is likely that precipitation has increased by 0.5 to 1% per decade in the 20<sup>th</sup> century over most mid- and high latitudes of the N.H. continents.**

ESS15  
Prof. Jin-Yi Yu


## Precipitation Changes (AR5)

Observed change in annual precipitation over land  
1951–2010




- Confidence in precipitation change averaged over global land areas since 1901 is low prior to 1951 and medium afterwards.
- Averaged over the mid-latitude land areas of the Northern Hemisphere, precipitation has likely increased since 1901 (medium confidence before and high confidence after 1951).
- For other latitudes area-averaged long-term positive or negative trends have low confidence


## Global Sea Level Has Risen



Tidal gauge data show that global average sea level rose between 10 and 20 cm during the 20th.




## Global Sea Level Change (AR5)



- Over the period 1901–2010, global mean sea level rose by 0.19 [0.17 to 0.21] m.
- The rate of sea level rise since the mid-19th century has been larger than the mean rate during the previous two millennia (high confidence).


## Changes in Cryosphere (AR5)



- Over the last two decades, the Greenland and Antarctic ice sheets have been losing mass (high confidence).
- Glaciers have continued to shrink almost worldwide (high confidence).
- Northern Hemisphere spring snow cover has continued to decrease in extent (high confidence).
- There is high confidence that there are strong regional differences in the trend in Antarctic sea ice extent, with a very likely increase in total extent.

## Summary from TAR

(a) Temperature Indicators



- Likelihood: {
- \*\*\* Virtually certain (probability > 99%)
  - \*\* Very likely (probability ≥ 90% but < 99%)
  - \* Likely (probability > 66% but < 90%)
  - ? Medium likelihood (probability > 33% but ≤ 66%)

in-Yi Yu



## Changes in Extreme Weather/Climate Events (AR5)

**Table SP4.1.1** Extreme weather and climate events: Global-scale assessment of recent observed changes, human contribution to the changes, and projected further changes for the early (2016–2035) and late (2081–2100) 21st century. Bold indicates where the AR5 (black) provides a revised<sup>1</sup> global-scale assessment from the SREX (blue) or AR4 (red). Projections for early 21st century were not provided in previous assessment reports. Projections in the AR5 are relative to the reference period of 1986–2005, and use the new Representative Concentration Pathway (RCP) scenarios (see Box SP4.1.1) unless otherwise specified. See the Glossary for definitions of extreme weather and climate events.

Phenomenon and direction of trend	Assessment that changes occurred typically since 1950 (unless otherwise indicated)	Assessment of a human contribution to observed changes	Likelihood of further changes			
			Early 21st century		Late 21st century	
Warmer and/or fewer cold days and nights over most land areas	Very likely	Very likely	Very likely	Very likely	Very likely	Very likely
Warmer and/or more frequent hot days and nights over most land areas	Very likely	Very likely	Very likely	Very likely	Very likely	Very likely
Warm spells/hot waves. Frequency and/or duration increases over most land areas	Medium confidence on a global scale. Likely in large parts of Europe, Asia and Australia.	Very likely	Not formally assessed	Very likely	Very likely	Very likely
Heavy precipitation events. Increase in the frequency, intensity, and/or amount of heavy precipitation	Likely more land areas with increases than decreases	Medium confidence	Likely over many land areas	Very likely	Very likely over most of the mid-latitude land masses and low-latitude tropical regions	Very likely
Increase in intensity and/or duration of drought	Low confidence on a global scale. Likely changes in some regions	Medium confidence	Low confidence	Likely (medium confidence) on a regional to global scale	More likely than not in the Western North Pacific and North Atlantic	Likely
Increase in intense tropical cyclone activity	Low confidence. Long-term (centennial) changes. Likely since 1970	Low confidence	Low confidence	More likely than not in the Western North Pacific and North Atlantic	More likely than not in some basins	Likely
Increased incidence and/or magnitude of extreme high sea level	Likely (since 1970)	Very likely	Very likely	Very likely	Very likely	Very likely

<sup>1</sup> The most comprehensive assessment to date, based on a different set of criteria, for some climate variables, different aspects have been assessed, and the revised guidance reduces uncertainties. The availability of new information, improved scientific understanding, continued analyses of data and models, and specific differences in methodologies applied in the assessed studies, all contribute to revised assessment findings.

<sup>2</sup> Attribution based on available case studies. It is likely that human influence has more than doubled the probability of occurrence of some observed heat waves in some locations.

<sup>3</sup> Model projections have been increased in the duration, intensity and spatial extent of heat waves and warm spells.

<sup>4</sup> In most estimates, confidence in trends is not higher than medium except in North America and Europe where there have been likely increases in either the frequency or intensity of heavy precipitation with some small and/or regional variation. It is very likely that there have been increases in recent North America.

<sup>5</sup> The frequency and intensity of drought has likely increased in the Mediterranean and West Africa, and likely decreased in central North America and north-west Australia.

<sup>6</sup> AR5 assessed the new affected drought.

<sup>7</sup> SREX assesses medium confidence that anthropogenic influence had contributed to some changes in the drought patterns observed in the second half of the 20th century, based on its attributed impact on precipitation and temperature changes. SREX assessed the attribution of changes in droughts to the new affected drought.

<sup>8</sup> There is a low confidence in projected changes in all metrics.

<sup>9</sup> Regional to global-scale projected decreases in total number and increased agricultural drought are likely (medium confidence) in presently dry regions by the end of the century under the RCP4.5 scenario. Substructure changes in the Mediterranean, Southeast US and southern European regions is consistent with projected changes in higher evaporation and increased surface temperatures, as there is high confidence in deep soil drying in these regions by the end of the century under the RCP4.5 scenario.

<sup>10</sup> There is medium confidence that a reduction in intense tropical cyclone activity is likely (medium confidence) in the North Atlantic basin and the Indian Ocean basin by the end of the century under the RCP4.5 scenario.

<sup>11</sup> Based on expert judgement and assessment of projections which use an SRES A1B or similar scenario.

<sup>12</sup> Attribution is based on the strong relationship between observed drought occurrence and sea level rise.

<sup>13</sup> There is high confidence that this increase in extreme high sea level will primarily be the result of an increase in mean sea level. There is low confidence in regional-specific projections of storminess and associated storm surge.

<sup>14</sup> SREX assessed to be very likely that mean sea level will contribute to future increased trends in extreme coastal high water levels.

## Major Conclusions in SPM

1. An increasing body of observations gives a collective picture of a warming world and other changes in the climate system.
2. Emissions of greenhouse gases and aerosols due to human activities continue to alter the atmosphere in ways that are expected to affect the climate.
3. Confidence in the ability of models to project future climate has increased.
4. There is new and stronger evidence that most of the warming observed over the last 50 years is attributable to human activities.
5. Human influences will continue to change atmospheric composition throughout the 21st century.
6. Global average temperature and sea level are projected to rise under all IPCC SRES scenarios.
7. Anthropogenic climate change will persist for many centuries.
8. Further action is required to address remaining gaps in information and understanding.


ESS15  
Prof. Jin-Yi Yu

## Emissions of Greenhouse Gases and Aerosols

- Concentrations of atmospheric greenhouse gases and their radiative forcing have continued to increase as a result of human activities.
- Anthropogenic aerosols are short-lived and mostly produce negative radiative forcing.

ESS15  
Prof. Jin-Yi Yu


## Radiative Forcing



$$\text{radiative forcing of the aerosols} = 15 - 20 = -5 \text{ w/m}^2$$

- Radiative forcing is a measure of the influence a factor has in altering the balance of incoming and outgoing energy in the Earth-atmosphere system, and is an index of the importance of the factor as a potential climate change mechanism. It is expressed in Watts per square meter ( $\text{Wm}^{-2}$ ).

ESS15  
Prof. Jin-Yi Yu



## Increase of CO<sub>2</sub> Concentration

- ❑ The atmospheric concentration of carbon dioxide (CO<sub>2</sub>) has increased by 31% since 1750. The present CO<sub>2</sub> concentration has not been exceeded during the past 420,000 years and likely not during the past 20 million years. The current rate of increase is unprecedented during at least the past 20,000 years.
  - ❑ About three-quarters of the anthropogenic emissions of CO<sub>2</sub> to the atmosphere during the past 20 years is due to fossil fuel burning. The rest is predominantly due to land-use change, especially deforestation.
  - ❑ Currently the ocean and the land together are taking up about half of the anthropogenic CO<sub>2</sub> emissions. On land, the uptake of anthropogenic CO<sub>2</sub> very likely exceeded the release of CO<sub>2</sub> by deforestation during the 1990s.
- ESS15 Prof. Jin-Yi Yu

## The Missing Carbon Sink

**Human Inject 6 Gigaton of Carbon Into the Atmosphere (per year)**

**3 Gigaton Remains in the Atmosphere**


**1.5 Gigaton Is Absorbed By the Ocean**

**Where Is he Missing 1.5 Gigaton (Sink)?**


*(Is this missing sink going to be saturated soon?)*

ESS15 Prof. Jin-Yi Yu

Variations in atmospheric CO<sub>2</sub> concentrations on different time-scales



## Ocean Acidification



**Figure SPM.4 |** Multiple observed indicators of a changing global carbon cycle: (a) atmospheric concentrations of carbon dioxide ( $\text{CO}_2$ ) from Mauna Loa ( $19^\circ 32' \text{N}$ ,  $155^\circ 34' \text{W}$  – red) and South Pole ( $89^\circ 59' \text{S}$ ,  $24^\circ 48' \text{W}$  – black) since 1958; (b) partial pressure of dissolved  $\text{CO}_2$  at the ocean surface (blue curves) and in situ pH (green curves), a measure of the acidity of ocean water. Measurements are from three stations from the Atlantic ( $29^\circ 10' \text{N}$ ,  $15^\circ 30' \text{W}$  – dark blue/dark green;  $31^\circ 40' \text{N}$ ,  $64^\circ 10' \text{W}$  – blue/green) and the Pacific Oceans ( $22^\circ 45' \text{N}$ ,  $158^\circ 00' \text{W}$  – light blue/light green). Full details of the datasets shown here are provided in the underlying report and the Technical Summary Supplementary Material (Figures 2.1 and 3.18; Figure TS.5)

- The ocean has absorbed about 30% of the emitted anthropogenic carbon dioxide, causing ocean acidification.
- Ocean acidification is quantified by decreases in pH. The pH of ocean surface water has decreased by 0.1 since the beginning of the industrial era (high confidence).

## Increase of $\text{N}_2\text{O}$ Concentration

- ❑ The atmospheric concentration of nitrous oxide ( $\text{N}_2\text{O}$ ) has increased by 46 ppb (17%) since 1750 and continues to increase. The present  $\text{N}_2\text{O}$  concentration has not been exceeded during at least the past thousand years.
- ❑ About a third of current  $\text{N}_2\text{O}$  emissions are anthropogenic (e.g., agricultural soils, cattle feed lots and chemical industry).



## Increase of Methane Concentration

- ❑ The atmospheric concentration of methane ( $\text{CH}_4$ ) has increased by 151% (1060 ppb<sub>9</sub>) since 1750 and continues to increase.
- ❑ The present  $\text{CH}_4$  concentration has not been exceeded during the past 420,000 years.
- ❑ Slightly more than half of current  $\text{CH}_4$  emissions are anthropogenic (e.g., use of fossil fuels, cattle, rice agriculture and landfills).



## Halocarbon Concentration

- ❑ Since 1995, the atmospheric concentrations of many of those halocarbon gases that are both ozone-depleting and greenhouse gases (e.g.,  $\text{CFCl}_3$  and  $\text{CF}_2\text{Cl}_2$ ), are either increasing more slowly or decreasing, both in response to reduced emissions under the regulations of the Montreal Protocol and its Amendments.
- ❑ Their substitute compounds (e.g.,  $\text{CHF}_2\text{Cl}$  and  $\text{CF}_3\text{CH}_2\text{F}$ ) and some other synthetic compounds (e.g., perfluorocarbons (PFCs) and sulphur hexafluoride ( $\text{SF}_6$ )) are also greenhouse gases, and their concentrations are currently increasing.



## Ozone Concentration

- ❑ The observed depletion of the stratospheric ozone (O<sub>3</sub>) layer from 1979 to 2000 is estimated to have caused a negative radiative forcing ( $-0.15 \text{ Wm}^{-2}$ ).
- ❑ The total amount of O<sub>3</sub> in the troposphere is estimated to have increased by 36% since 1750, due primarily to anthropogenic emissions of several O<sub>3</sub>-forming gases. This corresponds to a positive radiative forcing of  $0.35 \text{ Wm}^{-2}$ .





## Anthropogenic Aerosols

- ❑ The major sources of anthropogenic aerosols are fossil fuel and biomass burning. These sources are also linked to degradation of air quality and acid deposition.
- ❑ In general, the direct radiative forcing of aerosols is negative (except for black carbon fossil).
- ❑ There is much less confidence in the ability to quantify the total aerosol direct effect, and its evolution over time.
- ❑ Aerosols also vary considerably by region and respond quickly to changes in emissions.
- ❑ In addition to their direct radiative forcing, aerosols have an indirect radiative forcing through their effects on clouds.
- ❑ There is now more evidence for this indirect effect, which is negative, although of very uncertain magnitude.



## Precipitations

“Precipitation is any liquid or solid water particle that falls from the atmosphere and reaches the ground.”



# Cloud Seeding



- ❑ The objective is to convert some of the supercooled droplets in a cool clouds to ice and cause precipitation by the Bergeron process.
- ❑ Two primary methods are used to trigger the precipitation process.
- ❑ Dry ice is used to lower cloud temperature to a freezing point in order to stimulate ice crystal production leading to the Bergeron process.
- ❑ Silver iodide initiates the Bergeron process by directly acting as freezing nuclei.
- ❑ Under ideal conditions, seeding may enhance precipitation by about 10%.

ESS15  
Prof. Jin-Yi Yu

# Indirect Effects of Anthropogenic Aerosols


Effect	Cloud Types Affected	Process	Sign of Change in TOA Radiation	Potential Magnitude	Scientific Understanding
Cloud albedo effect	All clouds	For the same cloud water or ice content more but smaller cloud particles reflect more solar radiation	Negative	Medium	Low
Cloud lifetime effect	All clouds	Smaller cloud particles decrease the precipitation efficiency thereby presumably prolonging cloud lifetime	Negative	Medium	Very low
Semi-direct effect	All clouds	Absorption of solar radiation by absorbing aerosols affects static stability and the surface energy budget, and may lead to an evaporation of cloud particles	Positive or negative	Small	Very low
Glaciation indirect effect	Mixed-phase clouds	An increase in IN increases the precipitation efficiency	Positive	Medium	Very low
Thermodynamic effect	Mixed-phase clouds	Smaller cloud droplets delay freezing causing super-cooled clouds to extend to colder temperatures	Positive or negative	Medium	Very low

- ❑ The indirect effect of anthropogenic aerosols is probably negative, although of very uncertain magnitude.

ESS15  
Prof. Jin-Yi Yu


# Radiative Forcing (TAR)

Global Mean Radiative Forcing of Climate for year 2000 relative to 1750



- ❑ Natural and anthropogenic substances and processes that alter the Earth's energy budget are drivers of climate change.
- ❑ Radiative forcing (RF) quantifies the change in energy fluxes caused by changes in these drivers for 2011 relative to 1750.
- ❑ Positive RF leads to surface warming, negative RF leads to surface cooling.


# Radiative Forcing (AR5)



- ❑ The total anthropogenic RF for 2011 relative to 1750 is 2.29 W m<sup>-2</sup>, and it has increased more rapidly since 1970 than during prior decades.
- ❑ The total natural RF from solar irradiance changes and stratospheric volcanic aerosols made only a small contribution to the net radiative forcing throughout the last century, except for brief periods after large volcanic eruptions.

## Natural Factors of Radiative Forcing

- ❑ Since the late 1970s, satellite instruments have observed small oscillations due to the 11-year solar cycle. Mechanisms for the amplification of solar effects on climate have been proposed, but currently lack a rigorous theoretical or observational basis.
- ❑ Stratospheric aerosols from explosive volcanic eruptions lead to negative forcing, which lasts a few years. Several major eruptions occurred in the periods 1880 to 1920 and 1960 to 1991.
- ❑ The combined change in radiative forcing of the two major natural factors (solar variation and volcanic aerosols) is estimated to be negative for the past two, and possibly the past four, decades.





## Major Conclusions in SPM

1. An increasing body of observations gives a collective picture of a warming world and other changes in the climate system.
2. Emissions of greenhouse gases and aerosols due to human activities continue to alter the atmosphere in ways that are expected to affect the climate.
3. Confidence in the ability of models to project future climate has increased.
4. There is new and stronger evidence that most of the warming observed over the last 50 years is attributable to human activities.
5. Human influences will continue to change atmospheric composition throughout the 21st century.
6. Global average temperature and sea level are projected to rise under all IPCC SRES scenarios.
7. Anthropogenic climate change will persist for many centuries.
8. Further action is required to address remaining gaps in information and understanding.



## More Confidence in Climate Models



## Evidence of Global Warming

- ❑ Detection and attribution studies consistently find evidence for an anthropogenic signal in the climate record of the last 35 to 50 years.
- ❑ Simulations of the response to natural forcings alone (i.e., the response to variability in solar irradiance and volcanic eruptions) do not explain the warming in the second half of the 20th century.
- ❑ However, they indicate that natural forcings may have contributed to the observed warming in the first half of the 20th century.
- ❑ Most of the observed warming over the last 50 years is likely to have been due to the increase in greenhouse gas concentrations.





## Major Conclusions in SPM



1. An increasing body of observations gives a collective picture of a warming world and other changes in the climate system.
2. Emissions of greenhouse gases and aerosols due to human activities continue to alter the atmosphere in ways that are expected to affect the climate.
3. Confidence in the ability of models to project future climate has increased.
4. There is new and stronger evidence that most of the warming observed over the last 50 years is attributable to human activities.
5. Human influences will continue to change atmospheric composition throughout the 21st century.
6. Global average temperature and sea level are projected to rise under all IPCC SRES scenarios.
7. Anthropogenic climate change will persist for many centuries.
8. Further action is required to address remaining gaps in information and understanding.





## IPCC Special Report on Emission Scenarios (SRES)




## Projected CO<sub>2</sub> Emissions



## Projected CO<sub>2</sub> Concentration



## Projected Global Temperature (TAR)



□ The globally averaged surface temperature is projected to increase by 1.4 to 5.8°C over the period 1990 to 2100.

## Global Distribution of The Warming





Figure 20 The annual mean change of the temperature (colour shading) and its range (isolines) (Unit: °C) for the SRES scenario A2 (upper panel) and the SRES scenario B2 (lower panel). Both SRES scenarios show the period 2071 to 2100 relative to the period 1961 to 1990 and were performed by OAGCMs. [Based on Figures 9.10d and 9.10e]

It is very likely that nearly all land areas will warm more rapidly than the global average, particularly those at northern high latitudes in the cold season.


Prof. Jin-Yi Yu

## Vertical Distribution of Warming (AR4)



Prof. Jin-Yi Yu

## Projected Temperature Extreme (AR4)




- It is *very likely* that heat waves will be more intense, more frequent and longer lasting in a future warmer climate. Cold episodes are projected to decrease significantly.

- Almost everywhere, daily minimum temperatures are projected to increase faster than daily maximum temperatures, leading to a decrease in diurnal temperature range.

Prof. Jin-Yi Yu



## Future Projections (AR5)



- ❑ Global surface temperature change for the end of the 21st century is likely to exceed 1.5°C relative to 1850 to 1900 for all RCP scenarios except RCP2.6.
- ❑ It is very likely that the Arctic sea ice cover will continue to shrink and thin and that Northern Hemisphere spring snow cover will decrease during the 21st century as global mean surface temperature rises. Global glacier volume will further decrease.
- ❑ Earth System Models project a global increase in ocean acidification for all RCP scenarios.
- ❑ Most aspects of climate change will persist for many centuries even if emissions of CO<sub>2</sub> are stopped.


ESS15  
Prof. Jin-Yi Yu

## Projected Precipitation (TAR)

- ❑ Based on global model simulations and for a wide range of scenarios, global average water vapor concentration and precipitation are projected to increase during the 21<sup>st</sup> century.
- ❑ By the second half of the 21st century, it is likely that precipitation will have increased over northern mid- to high latitudes and Antarctica in winter.
- ❑ At low latitudes there are both regional increases and decreases over land areas.
- ❑ Larger year to year variations in precipitation are very likely over most areas where an increase in mean precipitation is projected.

ESS15  
Prof. Jin-Yi Yu


## Projected Changes in Cloud Cover (AR4)



Prof. Jin-Yi Yu

## Projected Precipitation Changes (AR4)


### a) Precipitation



- Precipitation generally increases in the areas of regional tropical precipitation maxima (such as the monsoon regimes) and over the tropical Pacific.
- Precipitation generally decreases in the subtropics and increases at high latitudes → due to the poleward shift of the storm track → due to the expansion of the Hadley circulation.


Prof. Jin-Yi Yu

## Properties of the Three Cells



ESS15  
Prof. Jin-Yi Yu

## Projected Precipitation Extreme (AR4)



- Intensity of precipitation events is projected to increase, particularly in tropical and high latitude areas that experience increases in mean precipitation.

- The number of dry days increases between precipitation events in the subtropics and lower mid-latitudes

- There is a tendency for drying of the mid-continental areas during summer, indicating a greater risk of droughts in those regions.


Prof. Jin-Yi Yu

## Projected Monsoon Variability (TAR)

- It is likely that warming associated with increasing greenhouse gas concentrations will cause an increase of Asian summer monsoon precipitation variability.
- Changes in monsoon mean duration and strength depend on the details of emission scenario.

ESS15  
Prof. Jin-Yi Yu

## Projected Monsoon Precipitation Changes (AR4)



Weaker monsoon circulation  
+ more water vapor available  
= increase monsoon rainfalls

West Africa Monsoon  
(increase)

Prof. Jin-Yi Yu

## Projected Hurricane Activities (AR4)

- Most recent published modelling studies projected a decrease in the overall number of storms.
- Although less confidence, studies projected decrease of relatively weak storms in most basins, with an increase in the numbers of the most intense tropical cyclones.




## Projected Extratropical Storms (AR4)

- For a future warmer climate, a poleward shift of storm tracks in both hemispheres that is particularly evident in the SH, with greater storm activity at higher latitudes.
- A future tendency for more intense extratropical storms, although the number of storms could be less.




## Projected Thermohaline Circulation (TAR)


- ❑ Most models show weakening of the ocean thermohaline circulation which leads to a reduction of the heat transport into high latitudes of the Northern Hemisphere.
- ❑ The current projections using climate models do not exhibit a complete shut-down of the thermohaline circulation by 2100.
- ❑ Beyond 2100, the thermohaline circulation could completely, and possibly irreversibly, shut-down in either hemisphere if the change in radiative forcing is large enough and applied long enough. scenarios.



## Projected Change in Atlantic Meridional Overturning Circulation (MOC) / AR4



- The MOC is an indicator of ocean circulation changes in response to global warming.
- The MOC is projected to slow down in the future.
- It is due to the warming and increased precipitation at higher latitudes.
- The weakened MOC will help to reduce global warming at higher latitude, because less heat will be transported there.




## Projected Snow/Ice Cover (TAR)


- ❑ Northern Hemisphere snow cover and sea-ice extent are projected to decrease further.
- ❑ Glaciers and ice caps are projected to continue their widespread retreat during the 21st century.
- ❑ The Antarctic ice sheet is likely to gain mass because of greater precipitation, while the Greenland ice sheet is likely to lose mass because the increase in runoff will exceed the precipitation increase.




## Projected Sea-Level Change (TAR)



- ❑ Global mean sea level is projected to rise by 0.09 to 0.88 metres between 1990 and 2100, for the full range of SRES scenarios.



## Global Warming and Sea-Level Change



## Positive Climate-Carbon Cycle Feedback

- Future climate change would reduce the efficiency of the Earth system (land and ocean) to absorb anthropogenic CO<sub>2</sub>.
- As a result, an increasingly large fraction of anthropogenic CO<sub>2</sub> would stay airborne in the atmosphere under a warmer climate.



Global mean sea level is projected to rise in the future warming world. The rise is contributed the most by this factor:

- a. Antarctic ice sheet melting
- b. Greenland ice sheet melting
- c. Glacier and ice cap melting
- d. thermal expansion



ESS15  
Prof. Jin-Yi Yu