


Lecture 9: Climate Sensitivity and Feedback Mechanisms

- ❑ Basic radiative feedbacks (Plank, Water Vapor, Lapse-Rate Feedbacks)
- ❑ Ice-albedo & Vegetation-Climate feedback
- ❑ Cloud feedback
- ❑ Biogeochemical feedbacks (skipped)
- ❑ Dynamical feedbacks and meridional energy transport (skipped)
- ❑ Longwave and evaporation feedbacks in the surface energy balance (skipped)



Climate Sensitivity and Feedback



(from *Earth's Climate: Past and Future*)



ESS200
Prof. Jin-Yi Yu

Definition and Mathematic Form

direct impact

$$\frac{dT_s}{dS_0} = \frac{\partial T_s}{\partial S_0} + \sum_{j=1}^N \frac{\partial T_s}{\partial y_j} \frac{dy_j}{dS_0}$$


□ Climate Sensitivity: the relationship between the measure of forcing and the magnitude of the climate change response.

□ Climate Forcing (S_0): Climate forcing is a change to the climate system that can be expected to change the climate.

□ Feedback Mechanism: a process that changes the sensitivity of the climate response.




Direct Impact and Feedback Process



Longwave Radiation (Plank) Feedback

$$E = \sigma T^4$$



- The outgoing longwave radiation emitted by the Earth depends on surface temperature, due to the Stefan-Boltzmann Law: $F = \sigma(T_s)^4$.
 - warmer the global temperature
 - larger outgoing longwave radiation been emitted by the Earth
 - reduces net energy heating to the Earth system
 - cools down the global temperature
 - a negative feedback



Stefan-Boltzmann Law

$$E = \sigma T^4$$

E = radiation emitted in W/m²


$\sigma = 5.67 \times 10^{-8}$ W/m² * K *sec

T = temperate (K ← *Kelvin degree*)

- ❑ The single factor that determines how much energy is emitted by a blackbody is its temperature.
- ❑ The intensity of energy radiated by a blackbody increases according to the fourth power of its absolute temperature.
- ❑ This relationship is called the Stefan-Boltzmann Law.




Water Vapor Feedback



- ❑ **Mixing Ratio** = the dimensionless ratio of the mass of water vapor to the mass of dry air.
- ❑ **Saturated Mixing Ratio** tells you the maximum amount of water vapor an air parcel can carry.
- ❑ The saturated mixing ratio is a function of air temperature: the warmer the temperature the larger the saturated mixing ratio.
 - ➔ a warmer atmosphere can carry more water vapor
 - ➔ stronger greenhouse effect
 - ➔ amplify the initial warming
 - ➔ one of the most powerful positive feedback



Clausius–Clapeyron Relationship



- Saturation vapor pressure describes how much water vapor is needed to make the air saturated at any given temperature.
- Saturation vapor pressure depends primarily on the air temperature in the following way:

$$\frac{de_s}{dT} = \frac{L}{T(\alpha_v - \alpha_l)}$$

**The
Clausius-Clapeyron
Equation**



$$e_s \cong 6.11 \cdot \exp \left\{ \frac{L}{R_v} \left(\frac{1}{273} - \frac{1}{T} \right) \right\}$$

- Saturation pressure increases exponentially with air temperature.

L: latent heat of evaporation; α : specific volume of vapor and liquid



ESS200
Prof. Jin-Yi Yu

Water Vapor Feedback


$$\frac{de_s}{dT} = \frac{L}{T(\alpha_v - \alpha_l)}$$

$$\frac{dq^*}{q^*} = \frac{de_s}{e_s} = \left(\frac{L}{R_v T} \right) \frac{dT}{T}$$

$$(20\%) = (20) \times (1\%)$$

- ❑ A 1% change in temperature (about 3°C) is associated with about a 20% change in saturation specific humidity
 - 1 °C change in temperature corresponds to a 7% change in saturation specific humidity.
- ❑ It is observed that the relative humidity of the atmosphere, which is the ratio of the actual to the saturation humidity, tends to remain constant, even when the air temperature goes through large seasonal variations in middle to high latitudes.
- ❑ In this case, we find that the terrestrial radiation emitted from the planet increases much less rapidly with temperature than would be indicated by the Stefan–Boltzmann relationship.
- ❑ An assumption that the relative humidity remains approximately constant when the climate warms or cools has generally been shown to be an excellent approximation.

Lapse-Rate Feedback



Global warming

- Decrease lapse rate
- Warm up middle troposphere
- Stronger terrestrial radiation
- Cooling climate
- A negative feedback

- ❑ In the tropics, the lapse rate is expected to decrease in response to the enhanced greenhouse effect, amplifying the warming in the upper troposphere and suppressing it at the surface.
- ❑ **This suppression causes a negative feedback on surface temperature.** Toward the poles, the reverse happens (a positive feedback), but the tropics tend to dominate, producing an overall negative feedback.
- ❑ Greenhouse gases and clouds cause the emission of thermal infrared radiation by Earth to originate in the middle troposphere, which is cooler than the surface.
- ❑ If we weaken the lapse rate, the surface and the emission temperatures become closer together, the greenhouse effect is weakened, and we should expect the surface temperature to cool.
- ❑ So if the lapse rate decreases with increasing surface temperature, that would be a negative feedback.




Figure 4.10: Schematic representation of positive and negative lapse-rate feedbacks.

http://www.climate.be/textbook/chapter4_node7.html



Temperature Changes due to Anthropogenic Warming





Figure 12.12 | CMIP5 multi-model changes in annual mean zonal mean temperature in the atmosphere and ocean relative to 1986–2005 for 2081–2100 under the RCP2.6 (left), RCP4.5 (centre) and RCP8.5 (right) forcing scenarios. Hatching indicates regions where the multi-model mean change is less than one standard deviation of internal variability. Stippling indicates regions where the multi-model change mean is greater than two standard deviations of internal variability and where at least 90% of the models agree on the sign of change (see Box 12.1).

(from IPCC AR5)

Snow/Ice Albedo Feedback



- ❑ The snow/ice albedo feedback is associated with the higher albedo of ice and snow than all other surface covering.
- ❑ This positive feedback has often been offered as one possible explanation for how the very different conditions of the ice ages could have been maintained.

TABLE 2-1 Average Albedo Range of Earth's Surfaces

Surface	Albedo range (percent)
Fresh snow or ice	60–90%
Old, melting snow	40–70
Clouds	40–90
Desert sand	30–50
Soil	5–30
Tundra	15–35
Grasslands	18–25
Forest	5–20
Water	5–10


Adapted from W. D. Sellers, Physical Climatology (Chicago: University of Chicago Press, 1965), and from R. G. Barry and R. J. Chorley, Atmosphere, Weather, and Climate, 4th ed. (New York: Methuen, 1982).

(from *Earth's Climate: Past and Future*)



ESS200
Prof. Jin-Yi Yu

Vegetation-Climate Feedbacks



(from *Earth' Climate: Past and Future*)



Cloud Feedback

Cloud Radiative Forcing as Estimated from Satellite Measurements

	Average	Cloud-free	Cloud forcing
OLR	234	266	+31
Absorbed solar radiation	239	288	-48
Net radiation	+5	+22	-17
Albedo	30%	15%	+15%

Radiative flux densities are given in W m^{-2} and albedo in percent. [From Harrison *et al.* (1990), © American Geophysical Union.]

- ❑ Clouds affect both solar radiation and terrestrial (longwave) radiation.
- ❑ Typically, clouds increase albedo → a cooling effect (negative feedback)
clouds reduce outgoing longwave radiation → a heating effect (positive feedback)
- ❑ The net effect of clouds on climate depends cloud types and their optical properties, the insolation, and the characteristics of the underlying surface.
- ❑ In general, high clouds tend to produce a heating (positive) feedback. Low clouds tend to produce a cooling (negative) feedback.
- ❑ Clouds double the albedo of Earth from 15% to 30% and reduce the longwave emission by about 30 W m^{-2} .
- ❑ The effect of clouds on the global net radiative energy flux into the planet is a reduction of about 20 W m^{-2} .