


# Lecture 3: ATMOSPHERE (Outline)



- Basic Structures and Dynamics
- General Circulation in the Troposphere
- General Circulation in the Stratosphere



**Energy (Heat)**




*The first law of thermodynamics*

**Air Temperature**



**Air Pressure**




**geostrophic balance**

**Air Motion**



# Air Pressure and Air Density




(from *Meteorology Today*)

- $\text{Weight} = \text{mass} \times \text{gravity}$
- $\text{Density} = \text{mass} / \text{volume}$
- $\text{Pressure} = \text{force} / \text{area}$   
 $= \text{weight} / \text{area}$



# Air Mass and Pressure




(from *Meteorology Today*)

- Atmospheric pressure tells you how much atmospheric mass is above a particular altitude.
- Atmospheric pressure decreases by about 10mb for every 100 meters increase in elevation.



# Hydrostatic Balance in the Vertical



(from *Climate System Modeling*)

□ vertical pressure force = gravitational force

$$-(dP) \times (dA) = \rho \times (dz) \times (dA) \times g$$


$$dP = -\rho g dz$$

$$dP/dz = -\rho g$$

***The hydrostatic balance !!***



# Hydrostatic Balance and Atmospheric Vertical Structure



(from *Meteorology Today*)

- Since  $P = \rho RT$  (the ideal gas law), the hydrostatic equation becomes:

$$dP = -P/RT \times g dz$$

- $dP/P = -g/RT \times dz$



- **$P = P_s \exp(-gz/RT)$**

- $P = P_s \exp(-z/H)$

- The atmospheric pressure decreases exponentially with height



# Temperature and Pressure





(from *Weather & Climate*)

- Hydrostatic balance tells us that the pressure decrease with height is determined by the temperature inside the vertical column.
- Pressure decreases faster in the cold-air column and slower in the warm-air column.
- Pressure drops more rapidly with height at high latitudes and lowers the height of the pressure surface.





# Thermal Energy to Kinetic Energy





# Pressure Gradient Force





(from Meteorology Today)

- ❑  $PG = (\text{pressure difference}) / \text{distance}$
- ❑ Pressure gradient force goes from high pressure to low pressure.
- ❑ Closely spaced isobars on a weather map indicate steep pressure gradient.




# Single-Cell Model: Explains Why There are Tropical Easterlies

## Without Earth Rotation



Coriolis Force

## With Earth Rotation




(Figures from *Understanding Weather & Climate* and *The Earth System*)



ESS220  
Prof. Jin-Yi Yu

# Coriolis Force



(from *The Earth System*)


- First, Point A rotates faster than Point B ( $U_A > U_B$ )
- $U_A > U_B$
- A northward motion starting at A will arrive to the east of B
- It looks like there is a “force” pushing the northward motion toward right
- This apparent force is called “Coriolis force”:

$$\text{Coriolis Force} = f V$$

$$\text{where } f = 2 * \Omega * \text{Sin}(\text{lat}) \text{ and } \Omega = 7.292 \times 10^{-5} \text{ rad s}^{-1}$$




# Another Kind of Coriolis Force



- ❑ The Coriolis force also causes the east-west wind to deflect to the right of its intent path in the Northern Hemisphere and to the left in the Southern Hemisphere.
- ❑ The deflections are caused by the centrifugal force associated with the east-west motion, and, therefore, related to rotation of the Earth, and are also considered as a kind of Coriolis force.
- ❑ Although the description of the deflection effect for north-south and east-west motions are very different, their mathematical expressions are the same.



# Coriolis Force Change with latitudes



(from *The Atmosphere*)




ESS220  
Prof. Jin-Yi Yu

# Coriolis Force

- ❑ Coriolis force causes the wind to deflect to the right of its intent path in the Northern Hemisphere and to the left in the Southern Hemisphere.
- ❑ The magnitude of Coriolis force depends on (1) the rotation of the Earth, (2) the speed of the moving object, and (3) its latitudinal location.
- ❑ The larger the speed (such as wind speed), the stronger the Coriolis force.
- ❑ The higher the latitude, the stronger the Coriolis force.
- ❑ The Coriolis force is zero at the equator.
- ❑ Coriolis force is one major factor that determine weather pattern.




# How Does Coriolis Force Affect Wind Motion?



(from *Weather & Climate*)



# Geostrophic Balance



□ By doing scale analysis, it has been shown that large-scale and synoptic-scale weather systems are in geostrophic balance.

□ Geostrophic winds always follow the constant pressure lines (isobar). Therefore, we can figure out flow motion by looking at the pressure distribution.







# Surface Friction

□ Friction Force =  $c * V$   
c = friction coefficient  
V = wind speed




# Frictional Effect on Surface Flow



- Surface friction force slows down the geostrophic flow.
- The flow turns into (out of) the low (high) press sides.
- Convergence (divergence) is produced with the flow.






(from *The Atmosphere*)



# Surface High and Low Pressure Systems




(from *The Atmosphere*)



ESS220  
Prof. Jin-Yi Yu


# Balance of Force in the Horizontal




Can happen in the tropics where the Coriolis force is small.

geostrophic balance

geostrophic balance plus frictional force



Upper Troposphere  
(free atmosphere)



Surface

(from *Weather & Climate*)





ESS220  
Prof. Jin-Yi Yu

# Surface Geostrophic Flow

## Cyclonic Flow


## Anticyclonic Flow




(figures from *Weather & Climate*)



# July



# Centrifugal Force



(from *The Atmosphere*)

□ The force that change the direction (but not the speed) of motion is called the centrifugal force.

□ Centrifugal Force =  $V^2 / R$ .  
V = wind speed  
R = the radius of the curvature






# Gradient Wind Balance

- The three-way balance of horizontal pressure gradient, Coriolis force, and the centrifugal force is call the *gradient wind balance*.
- The gradient wind is an excellent approximation to the actual wind observed above the Earth's surface, especially at the middle latitudes.



# Super- and Sub-Geostrophic Wind



(from *Meteorology: Understanding the Atmosphere*)

□ For high pressure system

→ gradient wind  $>$  geostrophic wind

→ supergeostrophic.


□ For low pressure system

→ gradient wind  $<$  geostrophic wind

→ subgeostrophic.




# Upper Tropospheric Flow Pattern



- Upper tropospheric flows are characterized by trough (low pressure; isobars dip southward) and ridge (high pressure; isobars bulge northward).
- The winds are in gradient wind balance at the bases of the trough and ridge and are slower and faster, respectively, than the geostrophic winds.
- Therefore, convergence and divergence are created at different parts of the flow patterns, which contribute to the development of the low and high systems.



# Developments of Low- and High-Pressure Centers





- **Dynamic Effects**: Combined curvature and jetstreak effects produce upper-level convergence on the west side of the trough to the north of the jetstreak, which add air mass into the vertical air column and tend to produce a surface high-pressure center. The same combined effects produce an upper-level divergence on the east side of the trough and favor the formation of a low-level low-pressure center.

- **Thermodynamic Effect**: heating → surface low pressure; cooling → surface high pressure.
- **Frictional Effect**: Surface friction will cause convergence into the surface low-pressure center after it is produced by upper-level dynamic effects, which adds air mass into the low center to “fill” and weaken the low center (increase the pressure)
- **Low Pressure**: The evolution of a low center depends on the relative strengths of the upper-level development and low-level friction damping.
- **High Pressure**: The development of a high center is controlled more by the convergence of surface cooling than by the upper-level dynamic effects. Surface friction again tends to destroy the surface high center.




# Single-Cell Model: Explains Why There are Tropical Easterlies

## Without Earth Rotation



Coriolis Force

## With Earth Rotation



(Figures from *Understanding Weather & Climate* and *The Earth System*)



ESS220  
Prof. Jin-Yi Yu

# Breakdown of the Single Cell → Three-Cell Model

□ Absolute angular momentum at **Equator** = Absolute angular momentum at **60°N**

□ The observed zonal velocity at the equator is  $u_{eq} = -5$  m/sec.

Therefore, the total velocity at the equator is  $U = \text{rotational velocity } (U_0 + u_{Eq})$

□ The zonal wind velocity at 60°N ( $u_{60N}$ ) can be determined by the following:

$$(U_0 + u_{Eq}) * a * \text{Cos}(0^\circ) = (U_{60N} + u_{60N}) * a * \text{Cos}(60^\circ)$$


$$(\Omega * a * \text{Cos}0^\circ - 5) * a * \text{Cos}0^\circ = (\Omega * a * \text{Cos}60^\circ + u_{60N}) * a * \text{Cos}(60^\circ)$$

$$u_{60N} = 687 \text{ m/sec !!!!}$$

**This high wind speed is not observed!**





# Properties of the Three Cells




# Atmospheric Circulation: Zonal-mean Views

## Single-Cell Model



## Three-Cell Model





(Figures from *Understanding Weather & Climate* and *The Earth System*)





# The Three Cells



Subtropical High

midlatitude Weather system

(Figures from *Understanding Weather & Climate* and *The Earth System*)



ESS220  
Prof. Jin-Yi Yu

# Thermally Direct/Indirect Cells


## ❑ Thermally Direct Cells (Hadley and Polar Cells)

Both cells have their rising branches over warm temperature zones and sinking branches over the cold temperature zone. Both cells directly convert thermal energy to kinetic energy.

## ❑ Thermally Indirect Cell (Ferrel Cell)

This cell rises over cold temperature zone and sinks over warm temperature zone. The cell is not driven by thermal forcing but driven by eddy (weather systems) forcing.





(from *The Earth System*)



ESS220  
Prof. Jin-Yi Yu

# Is the Three-Cell Model Realistic?

□ *Yes and No!*

*(Due to sea-land contrast and topography)*

*Yes:* the three-cell model explains reasonably well the surface wind distribution in the atmosphere.

*No:* the three-cell model can not explain the circulation pattern in the upper troposphere. (planetary wave motions are important here.)



# Semi-Permanent Pressure Cells

## □ **The Aleutian, Icelandic, and Tibetan lows**


- The oceanic (continental) lows achieve maximum strength during winter (summer) months
- The summertime Tibetan low is important to the east-Asia monsoon

## □ **Siberian, Hawaiian, and Bermuda-Azores highs**


- The oceanic (continental) highs achieve maximum strength during summer (winter) months




# January




# July



# Sinking Branches and Deserts



(a)



(b)




(c)

(from *Weather & Climate*)






# Global Distribution of Deserts



(from *Global Physical Climatology*)



# Upper Tropospheric Circulation



(from *Weather & Climate*)

- ❑ Only the Hadley Cell can be identified in the lower latitude part of the circulation.
- ❑ Circulation in most other latitudes are dominated by westerlies with wave patterns.
- ❑ Dominated by large-scale waver patterns (wave number 3 in the Northern hemisphere).



# Subtropical and Polar Jet Streams





FIG. 3.8 Winter (December–February) zonal mean wind components (knots), Northern Hemisphere, at (a) 140°E and (b) 0° longitude. (Redrawn from Crutcher, 1961.)

## □ Subtropical Jet

Located at the higher-latitude end of the Hadley Cell. The jet obtain its maximum wind speed (westerly) due the conservation of angular momentum.

## □ Polar Jet


Located at the thermal boundary between the tropical warm air and the polar cold air. The jet obtain its maximum wind speed (westerly) due the latitudinal thermal gradient (thermal wind relation).



(from *Atmospheric Circulation Systems*)



# Thermal Wind Relation



(from *Weather & Climate*)



# Thermal Wind Equation



$$\frac{\partial U}{\partial z} \propto - \frac{\partial T}{\partial y}$$

- The vertical shear of zonal wind is related to the latitudinal gradient of temperature.
- Jet streams usually are formed above baroclinic zone (such as the polar front).



# Jet Streams Near the Western US

*Pineapple Express*



- ❑ Both the polar and subtropical jet streams can affect weather and climate in the western US (such as California).
- ❑ El Nino can affect western US climate by changing the locations and strengths of these two jet streams.

(from Riehl (1962), Palmen and Newton (1969))



ESS220  
Prof. Jin-Yi Yu

# Parameters Determining Mid-latitude Weather


- Temperature differences between the equator and poles
- The rate of rotation of the Earth.



# Rotating Annulus Experiment

Cooling  
Outside

Heating  
Inside




(from “*Is The Temperature Rising?*”)



ESS220  
Prof. Jin-Yi Yu



# New Understanding of Cyclone after WWII



**Carl Gustav Rossby (1898-1957)**

- ❑ Carl Rossby mathematically expressed relationships between mid-latitude cyclones and the upper air during WWII.
- ❑ Mid-latitude cyclones are a large-scale waves (now called Rossby waves) that grow from the “baroclinic” instability associated with the north-south temperature differences in middle latitudes.



# Polar Front Theory




Vilhelm Bjerknes (1862-1951)

- *Bjerknes*, the founder of the Bergen school of meteorology, developed polar front theory during WWI to describe the formation, growth, and dissipation of mid-latitude cyclones.



# El Nino and Southern Oscillation


- ❑ Jacob Bjerknes was the first one to recognize that El Nino is not just an oceanic phenomenon (in his 1969 paper).
- ❑ In stead, he hypothesized that the warm waters of El Nino and the pressure seasaw of Walker's Southern Oscillation are part and parcel of the same phenomenon: the ENSO.
- ❑ Bjerknes's hypothesis of coupled atmosphere-ocean instability laid the foundation for ENSO research.



Jacob Bjerknes



(From *The Blue Planet*)



# How Cyclone Grows?

Potential Energy  $\rightarrow$  Available P. Energy

(cold/warm air moves south/north)

$$(V^*T^* > 0)$$


Available Energy  $\rightarrow$  Kinetic Energy

(cold/warm air moves down/up)


$$(W^*T^* > 0)$$



# Available Potential Energy



# Life Cycle of Mid-Latitude Cyclone




(from *Weather & Climate*)

- Cyclogenesis
- Mature Cyclone
- Occlusion




# Cold and Warm Fronts


## Mid-Latitude Cyclone



(From *Weather & Climate*)




cold front



warm front






ESS220  
Prof. Jin-Yi Yu



# Tropical Hurricane




- The hurricane is characterized by a strong thermally direct circulation with the rising of warm air near the center of the storm and the sinking of cooler air outside.

(from *Understanding Weather & Climate*)




# They Are the Same Things...



- ❑ **Hurricanes:** extreme tropical storms over Atlantic and eastern Pacific Oceans.
- ❑ **Typhoons:** extreme tropical storms over western Pacific Ocean.
- ❑ **Cyclones:** extreme tropical storms over Indian Ocean and Australia.




# Walker Circulation and Ocean Temperature



# East-West Circulation

(from Flohn (1971))




□ The east-west circulation in the atmosphere is related to the sea/land distribution on the Earth.




## Southern Oscillation: an atmospheric phenomenon

In 1910s, Walker found a connection between barometer readings at stations on the eastern and western sides of the Pacific (Tahiti and Darwin). He coined the term **Southern Oscillation** to dramatize the ups and downs in this east-west seesaw effect.




(from Rasmusson 1984)







Sir Gilbert Walker




# Walker Circulation and Ocean




# Scales of Motions in the Atmosphere



# Monsoon: Sea/Land-Related Circulation



Courtesy of Kevin G. Cannariato



- ❑ Monsoon (Arabic “season”)
- ❑ Monsoon is a climate feature that is characterized by the *seasonal reversal in surface winds*.
- ❑ The very different heat capacity of land and ocean surface is the key mechanism that produces monsoons.
- ❑ During summer seasons, land surface heats up faster than the ocean. Low pressure center is established over land while high pressure center is established over oceans. Winds blow from ocean to land and bring large amounts of water vapor to produce heavy precipitation over land: A rainy season.
- ❑ During winters, land surface cools down fast and sets up a high pressure center. Winds blow from land to ocean: a dry season.






# How Many Monsoons Worldwide?

North America Monsoon

Asian Monsoon



Australian Monsoon

South America Monsoon


Africa Monsoon

(figure from *Weather & Climate*)



ESS220  
Prof. Jin-Yi Yu

# Orbital-Scale Changes in Methane




(from *Earth's Climate: Past and Future*)

- ❑ The Vostok ice record shows a series of cyclic variations in methane concentration, ranging between 350 to 700 ppb (part per billion).
- ❑ Each CH<sub>4</sub> cycle takes about 23,000 years.
- ❑ This cycle length points to a likely connection with changes in orbital procession.
- ❑ The orbital procession dominates insolation changes at *lower latitudes*.



ESS220  
Prof. Jin-Yi Yu

# Trapping Gases in the Ice



- ❑ Air moves freely through snow and ice in the upper 15 m of an ice sheet.
- ❑ Flow is increasingly restricted below this level.
- ❑ Bubbles of old air are eventually sealed off completely in ice 50 to 100 m below the surface.




# Monsoon and Methane

- ❑ On the 23,000-year cycle, methane variations closely resemble the variations of monsoon strength.
- ❑ The peak values of methane match the expected peaks in monsoon intensity not only in timing but also in amplitude.
- ❑ This match suggests a close connection between CH<sub>4</sub> concentrations and the monsoon on the 23,000-year climate cycle.
- ❑ By why?



# Earth's Orbit and Its Variations




(from *The Earth System*)

- ❑ First, Earth spins around on its axis once every day → The *Tilt*.
- ❑ Second, Earth revolves around the Sun once a year → The shape of the *Orbit*.
- ❑ Both the tilt and the shape of the orbit have changed over time and produce three types of orbital variations:
  - (1) obliquity variations
  - (2) eccentricity variations
  - (3) precession of the spin axis.



ESS220  
Prof. Jin-Yi Yu

# Precession of Axis




(from *Earth's Climate: Past and Future*)

- ❑ There are two kinds of precession: (1) the precession of the spin axis and (2) the precession of the ellipse.
- ❑ Earth's wobbling motion is called the axial precession. It is caused by the gravitational pull of the Sun and Moon.
- ❑ Axial precession is a slow turning of Earth's axis of rotation through a circular path, with a full turn every 25,700 years.



ESS220  
Prof. Jin-Yi Yu

# Precession of Ellipse




- The precession of the ellipse is known as the elliptical shape of Earth's orbit rotates itself at a slower rate than the wobbling motion of the axial precession.

(from *Earth's Climate: Past and Future*)



ESS220  
Prof. Jin-Yi Yu

# Time Scales of Precession




(from *Earth's Climate: Past and Future*)

- The combined effects of these two precessions cause the solstices and equinoxes to move around Earth's orbit, completing one full 360° orbit around the Sun every 23,000 years.





# The Orbital Monsoon Hypothesis



(from *Earth's Climate: Past and Future*)

- ❑ The 23,000-year cycle of orbital procession increases (decreases) summer insolation and at the same time decreases (increases) winter insolation at low and middle latitudes.
- ❑ Departures from the modern seasonal cycle of solar radiation have driven stronger monsoon circulation in the past.
- ❑ Greater summer radiation intensified the wet summer monsoon.
- ❑ Decreased winter insolation intensified the dry winter monsoon.



# How Did Monsoon Affect Methane?


- Orbital procession affects solar radiation at low latitudes
  - solar radiation affects the strength of low-latitude monsoons
  - monsoon fluctuations changes the precipitation amounts in *Southeast Asia*
  - heavy rainfalls increase the amount of standing water in bogs
  - decaying vegetation used up any oxygen in the water and creates the oxygen-free conditions needed to generate methane
  - the extent of these boggy area must have expanded during wet monsoon maximum and shrunk during dry monsoon minimum.



# Seasonal Cycle of Rainfall

(from IRI)

Mean Jan GPCP Precipitation (79-03)





Prof. Jin-Yi Yu


# Computer Simulation of Eastern Pacific Warm Pool



Topography (meter)



# Sea/Land Breeze



- ❑ Sea/land breeze is also produced by the different heat capacity of land and ocean surface, similar to the monsoon phenomenon.
- ❑ However, sea/land breeze has much shorter timescale (day and night) and space scale (a coastal phenomenon) than monsoon (a seasonal and continental-scale phenomenon).

(figure from *The Earth System*)



# Santa Ana Wind

---



This is a picture of Fremont Canyon, located in the Santa Ana Mountains in Orange County. This canyon is known for its extremely high winds during Santa Ana wind events, where the winds can gust over 100 MPH during very strong Santa Ana wind events (picture from the Orange County Register).

## DEFINITION

**Strong warm and dry winds blow over the southern California from the Great Basin, with speeds exceed 25 knots (46 km/hr).**




# Generation Mechanism



(from NASA's Observatorium website)



10/10/2003



## Santa Ana Wind

The air is forced down the mountain slopes towards the Pacific coast

Dry, low humidity and hot, with sinking air temperature 40C (104F) near the coast


Often contribute to the spread of severely destructive wild fires in California






# Santa Ana Wind

Santa Ana winds on February 9, 2002  
NASA MISR observation



Santa Ana Guide ©1999 Channel Crossings Press



ESS220  
Prof. Jin-Yi Yu

# **Diurnal and Seasonal Variations**

---

## **Diurnal variation:**

**Stronger Santa Ana wind at night and weaker Santa Ana wind on the day.**

## **Seasonal Variation:**

**Occurs most frequently in winter (November to March).**




10/22/2007



# Temperatures in Stratosphere

stratosphere mesosphere




(from *Dynamic Meteorology*)




ESS220  
Prof. Jin-Yi Yu

# Ozone Distribution



**Fig. 12.13** Time-latitude section showing the seasonal variation of total ozone (Dobson units) based on total ozone mapping spectrometer (TOMS) data. Note the extratropical springtime maxima in both hemispheres and the minimum at 90°S centered on October 1. (After Bowman and Krueger, 1985.)



# Stratosphere: Circulation and Temperature

## Zonal Wind




Fig. 1.4. Schematic latitude-height section of zonal mean zonal wind ( $\text{m s}^{-1}$ ) for solstice conditions; W and E designate centers of westerly (from the west) and easterly (from the east) winds, respectively. (Courtesy of R. J. Reed.)

## Temperature





Fig. 1.3. Schematic latitude-height section of zonal mean temperatures ( $^{\circ}\text{C}$ ) for solstice conditions. Dashed lines indicate tropopause, stratopause, and mesopause levels. (Courtesy of R. J. Reed.)



# Circulation in Stratosphere




(from *Dynamic Meteorology*)



ESS220  
Prof. Jin-Yi Yu

# Zonal-Mean Circulation in the Stratosphere



summer

winter





# Ozone Production and Destruction

(from *The Earth System*)

## The Chapman Mechanism of Ozone Production and Destruction

Photodissociation  
(or photolysis)

**Reaction\***

**Rate**

1)	$O_2 + \text{UV photon} \rightarrow O + O$	} production	Slow
2)	$O + O_2 + M \rightarrow O_3 + M$		Fast
3)	$O_3 + \text{photon} \rightarrow O_2 + O$	} destruction	Fast
4)	$O + O_3 \rightarrow 2 O_2$		Slow


visible light

destroy O3 permanently



ESS220  
Prof. Jin-Yi Yu

# Ozone Distribution



(from *The Earth System*)

- ❑ The greatest production of ozone occurs in the tropics, where the solar UV flux is the highest.
- ❑ However, the general circulation in the stratosphere transport ozone-rich air from the tropical upper stratosphere to mid-to-high latitudes.
- ❑ Ozone column depths are highest during springtime at mid-to-high latitudes.
- ❑ Ozone column depths are the lowest over the equator.



# Climate Variations in Stratosphere

- ❑ Quasi-Biennial Oscillation (QBO)
- ❑ Sudden Warming: in Northern Pole
- ❑ Ozone Hole: in Southern Pole



# QBO

□ Quasi-Biennial Oscillation: Easterly and westerly winds alternate every other years (approximately) in the lower to middle parts of the tropical stratosphere.





Fig. 8.1. Time-height section of monthly mean zonal winds ( $\text{m s}^{-1}$ ) at equatorial stations (Jan. 1953–Aug. 1967: Canton Island,  $3^{\circ}\text{S}/172^{\circ}\text{W}$ ; Sept. 1967–Dec. 1975: Gan/Maldives Islands,  $1^{\circ}\text{S}/73^{\circ}\text{E}$ ; Jan. 1976–Apr. 1985: Singapore,  $1^{\circ}\text{N}/104^{\circ}\text{E}$ ). Isopleths are at  $10\text{-m s}^{-1}$  intervals. Note the alternating downward propagating westerly (W) and easterly (E) regimes. [From Naujokat (1986), with permission.]



# Why QBO?




**Fig. 8.7.** Schematic representation of the evolution of the mean flow in Plumb's analog of the QBO. Six stages of a complete cycle are shown. Double arrows show wave-driven accelerations and single arrows show viscously driven accelerations. Wavy lines indicate relative penetration of easterly and westerly waves. See text for details. [After Plumb (1984).]

- Kelvin Waves accelerate westerly.
- Rossby-Gravity Wave accelerate easterly.



# Sudden Warming



- ❑ Every other year or so the normal winter pattern of a cold polar stratosphere with a westerly vortex is interrupted in the middle winter.
- ❑ The polar vortex can completely disappear for a period of a few weeks.
- ❑ During the sudden warming period, the stratospheric temperatures can rise as much as  $40^{\circ}\text{K}$  in a few days!




# Why Sudden Warming?

- ❑ Planetary-scale waves propagating from the troposphere (produced by big mountains) into the stratosphere.
- ❑ Those waves interact with the polar vortex to break down the polar vortex.
- ❑ There are no big mountains in the Southern Hemisphere to produce planetary-scale waves.
- ❑ Less (?) sudden warming in the southern polar vortex.



# Antarctic Ozone Hole

Mean Total Ozone Over Antarctic in October



- ❑ The decrease in ozone near the South Pole is most striking near the spring time (October).
- ❑ During the rest of the year, ozone levels have remained close to normal in the region.

(from *The Earth System*)




ESS220  
Prof. Jin-Yi Yu



# The 1997 Ozone Hole

Total Ozone for Aug 1, 1997




GSFC/918



ESS220  
Prof. Jin-Yi Yu

# Why No Ozone Hole in Arctic?



(from WMO Report 2003)



ESS220  
Prof. Jin-Yi Yu

# Polar Stratospheric Clouds (PSCs)



(Sweden, January 2000; from NASA website)

- ❑ In winter the polar stratosphere is so cold ( $-80^{\circ}\text{C}$  or below) that certain trace atmospheric constituents can condense.
- ❑ These clouds are called “polar stratospheric clouds” (PSCs).
- ❑ The particles that form typically consist of a mixture of water and nitric acid ( $\text{HNO}_3$ ).
- ❑ The PSCs alter the chemistry of the lower stratosphere in two ways:
  - (1) by coupling between the odd nitrogen and chlorine cycles
  - (2) by providing surfaces on which heterogeneous reactions can occur.



# Ozone Hole Depletion

- ❑ Long Antarctic winter (May through September)
- ➔ The stratosphere is cold enough to form PSCs
- ➔ PSCs deplete odd nitrogen (NO)
- ➔ Help convert unreactive forms of chlorine (ClONO<sub>2</sub> and HCl) into more reactive forms (such as Cl<sub>2</sub>).
- ➔ The reactive chlorine remains bound to the surface of clouds particles.
- ➔ Sunlight returns in springtime (September)
- ➔ The sunlight releases reactive chlorine from the particle surface.
- ➔ The chlorine destroy ozone in October.
- ➔ Ozone hole appears.
- ➔ At the end of winter, the polar vortex breaks down.
- ➔ Allow fresh ozone and odd nitrogen to be brought in from low latitudes.
- ➔ The ozone hole recovers (disappears) until next October.

