




Mid-Latitude Cyclones

- Mid-latitude cyclones form along a boundary separating polar air from warmer air to the south.
- □ These cyclones are large-scale systems that typically travels eastward over greart distance and bring precipitations over wide areas.
- □ Lasting a week or more.











- polar front but may initiate elsewhere, such
- boundary separating colder polar easterlies
- A low pressure area forms and due to the counterclockwise flow (N.H.) colder air migrates equatorward behind a developing
- developing warm front (east of the system).
- association with converging winds of the ESS5 Prof. Jin-Yi Yu







Mature Cyclones


- U Well-developed fronts circulating about a deep low pressure center characterize a mature mid-latitude cyclone.
- Heavy precipitation stems from cumulus development in association with the cold front.
- Lighter precipitation is associated with stratus clouds of the warm front.
- □ Isobars close the low and are typically kinked in relation to the fronts due to steep temperature gradients.


ESS5 Prof. Jin-Yi Yu
































Vortocity and Divergence




- Decreasing vorticity in the zone between a trough and ridge leads to upper air convergence and sinking motions through the atmosphere, which supports surface high pressure areas.
- □ Increasing vorticity in the zone between a ridge and trough leads to upper air divergence and rising motions through the atmosphere, which supports surface low pressure areas.











- movement of the storm system is seen through a comparison of weather maps over a 24-hour period
- the low moves over the northern Great Lakes
- In the upper air, the trough has increased in amplitude and strength and become oriented northwest to southeast Isobars have closed about the low,






- April 17 Continual movement towards the northeast is apparent, although system movement has lessened
- The occlusion is now sweeping northeastward of the low, bringing snowfall to regions to the
- In the upper air, continued deepening is occurring in association with the more robust cutoff low



Steering of Mid-Latitude Cyclones

- □ The movement of surface systems can be predicted by the 500 mb pattern.
- The surface systems move in about the same direction as the 500 mb flow, at about 1/2 the speed.
- Upper-level winds are about twice as strong in winter than summer.
- This results in stronger pressure gradients (and winds), resulting in stronger and more rapidly moving surface cyclones.





<section-header>