


## Chapter 7: Precipitation Processes



From: Introduction to Tropical Meteorology, 1st Edition, Version 1.1.2,  
Produced by the [COMET® Program](#)  
Copyright 2007-2008, University Corporation for Atmospheric Research.



100% of 1477KB loaded.

From: Introduction to Tropical Meteorology, 1st Edition, Version 1.1.2,  
Produced by the [COMET® Program](#)  
Copyright 2007-2008, University Corporation for Atmospheric Research.



- Growth of Cloud Droplet
- Forms of Precipitations
- Cloud Seeding



# Precipitations

Water Vapor Saturated



Need Cloud Condensation Nuclei

Cloud Droplet formed around Cloud Condensation Nuclei




Need to fall down


Precipitation



Key:  
 $r$  = radius in micrometers  
 $n$  = number per liter  
 $V$  = terminal velocity in centimeters per second



# Terminal Velocity



- ❑ Terminal velocity is the constant speed that a falling object has when the gravity force and the drag force applied on the subject reach a balance.
- ❑ Terminal velocity depends on the size of the object: small objects fall slowly and large objectives fall quickly.



# Raindrops

- ❑ Rain droplets have to have large enough falling speed in order to overcome the updraft (that produces the rain) to fall to the ground.
- ❑ This means the rain droplets have to **GROW** to large enough sizes to become precipitation.





## How Raindrop Grows?

- ❑ Growth by Condensation (too small)
- ❑ Growth in Warm Clouds:
  - Collision-Coalescence Process
- ❑ Growth in Cool and Cold Clouds:
  - Bergeron Process




## Growth by Condensation

- ❑ Forms most cloud drops initially
- ❑ Growth limited to a radii of  $\sim 20 \mu\text{m}$   
Insufficient for precipitation




## Growth in Warm Clouds




- ❑ Most clouds formed in the Tropics, and many in the middle latitudes, are warm clouds (greater than  $0^\circ\text{C}$ ).
- ❑ The **Collision-coalescence** process generates precipitation.
- ❑ This process depends on the differing fall speeds of different-sized droplets.
- ❑ It begins with large collector drops which have high terminal velocities.






(a)



(b)



(c)

## Collision

- ❑ Collector drops collide with smaller drops.
- ❑ Due to compressed air beneath falling drop, there is an inverse relationship between collector drop size and collision efficiency.
- ❑ Collisions typically occur between a collector and fairly large cloud drops.
- ❑ Smaller drops are pushed aside.
- ❑ Collision is more effective for the droplets that are not very much smaller than the collect droplet.



## Coalescence

- ❑ When collisions occur, drops either bounce apart or coalesce into one larger drop.
- ❑ Coalescence efficiency is very high indicating that most collisions result in coalescence.
- ❑ Collision and coalescence together form the primary mechanism for precipitation in the tropics, where warm clouds dominate.



## Cool and Cold Clouds


- ❑ A portion of most mid-latitude clouds have temperatures below the melting point of ice.
- ❑ Cold clouds are referred to those have temperature below 0°C throughout and consist entirely of ice crystals, supercooled droplets, or a mixture of two.
- ❑ Cool clouds are referred to those have temperatures above 0°C in the lower reaches and subfreezing condition above.



## An Example of Cool and Cold Cloud




Cumulonimbus clouds contain both ice (top, fuzzy cloud margins), liquid drops (bottom, sharp margins) and a mix of ice and liquid (middle)



## Growth in Cool and Cold Clouds

- ❑ Cool month mid-latitude and high latitude clouds are classified as cool clouds as average temperatures are usually below freezing.
- ❑ Clouds may be composed of (1) Liquid water, (2) Supercooled water, and/or (3) Ice.
- ❑ Coexistence of ice and supercooled water is critical to the creation of cool cloud precipitation - the **Bergeron Process**.

## Bergeron Process



- ❑ Saturation vapor pressure of ice is less than that of supercooled water and water vapor.
- ❑ Saturation vapor pressure gradient allows rapid growth of ice crystals

## Riming and Aggregation

- ❑ Collisions between falling crystals and drops causes growth through *riming* and *aggregation*.
- ❑ *Riming* = liquid water freezing onto ice crystals producing rapid growth.
- ❑ *Aggregation* = the joining of multiple ice crystals through the bonding of surface water builds ice crystals to the point of overcoming updrafts
- ❑ Collision combined with riming and aggregation allow formation of precipitation within 1/2 hour of initial formation.





## Forms of Precipitation

- ❑ *Snow*
- ❑ *Rain*
- ❑ *Graupel and Hail*
- ❑ *Sleet*
- ❑ *Freezing Rain*




## Snow






- ❑ Snowflakes have a wide assortment of shapes and sizes depending on moisture content and temperature of the air.
- ❑ Snowfall distribution in North America is related to north-south alignment of mountain ranges and the presence of the Great Lakes.





## Rain

- Rainshowers: are episodic precipitation events associated with convective activity and cumulus clouds
- Raindrop shape:
  - As frictional drag increases, changes to a mushroom shape
  - Drops eventually flatten
  - Drops split when frictional drag overcomes the surface tension of water
  - Splitting ensures a maximum drop size of about 5 mm and the continuation of the collision-coalescence process




## Graupel and Hail

- **Graupel** are ice crystals that undergo extensive riming
  - Lose six sided shape and smooth out
  - Either falls to the ground or provides a nucleus for hail
- **Hail** are concentric layers of ice around graupel
  - Up and downdraft interactions in thunderstorms
  - Great Plains = highest frequency of hail events



## Hail Formation



Concentric layers of ice in hail indicate the cyclical hailstone formation process




## Hail Frequency in the U.S.






## Sleet and Freezing Rain



- **Sleet** begins as ice crystals which melt into rain through a mid-level inversion before solidifying in colder near surface air
- **Freezing Rain** forms similarly to sleet, but does not fully refreeze



## Twomey effect, 1974

How CCN (cloud condensation nuclei) from anthropogenic pollution may

increase the cloud lifetime and decrease the precipitation



## Cloud Seeding

- ❑ The objective is to convert some of the supercooled droplets in a cool clouds to ice and cause precipitation by the **Bergeron process**.
- ❑ Two primary methods are used to trigger the precipitation process.
- ❑ **Dry ice** is used to lower cloud temperature to a freezing point in order to stimulate ice crystal production leading to the Bergeron process.
- ❑ **Silver iodide** initiates the Bergeron process by directly acting as freezing nuclei.
- ❑ Under ideal conditions, seeding may enhance precipitation by about 10%.



## Measuring Precipitation

- Standard *raingages*
  - Sparse network



(4)



provide a record of precipitation amount/time of the event



### ❑ Raingage Measurement Errors


- Point estimates
  - Wide variations across small spaces
  - Overestimates
    - Wind, residual water

### ❑ Precipitation Measurement by Weather Radar

- Precipitation estimates
- Real-time information



## Doppler Radar Precipitation Estimates



## Measuring Snow

- Accumulated snow measured
  - Water equivalent of snow = 10 to 1 ratio
- Automated snow pillows
  - Convert weight to water equivalent

